arXiv:1208.1526v1 [math.GT] 7 Aug 2012

THE EQUIVALENCE OF HEEGAARD FLOER HOMOLOGY AND EMBEDDED CONTACT HOMOLOGY III: FROM HAT TO PLUS

VINCENT COLIN, PAOLO GHIGGINI, AND KO HONDA

ABSTRACT. Given a closed oriented 3-manifold M, we establish an isomorphism between the Heegaard Floer homology group $HF^+(-M)$ and the embedded contact homology group $ECH(M)$. Starting from an open book decomposition $(S,\hbar)$ of M, we construct a chain map Φ^+ from a Heegaard Floer chain complex associated to $(S,\hbar)$ to an embedded contact homology chain complex for a contact form supported by $(S,\hbar)$. The chain map Φ^+ commutes up to homotopy with the U-maps defined on both sides and reduces to the quasi-isomorphism Φ from [CGH2, CGH3] on subcomplexes defining the hat versions. Algebraic considerations then imply that the map Φ^+ is a quasi-isomorphism.

CONTENTS

1. Introduction 1
2. Heegaard Floer chain complexes 4
3. The geometric U-map 5
4. The cobordism W^+ 12
5. The chain map Φ^+ 18
6. Proof of Theorem 1.0.1 25

References 31

1. INTRODUCTION

This is the last paper in the series which proves the isomorphism between certain Heegaard Floer homology and embedded contact homology groups. References from [CGH2] (resp. [CGH3]) will be written as "Section I.x" (resp. "Section II.x") to mean "Section x" of [CGH2] (resp. [CGH3]), for example.

Let M be a closed oriented 3-manifold. Let $\widehat{HF}(M)$ and $HF^+(M)$ be the hat and plus versions of Heegaard Floer homology of M and let $\widehat{ECH}(M)$ and $ECH(M)$ be hat and usual versions of the embedded contact homology of M. As

---

Date: This version: August 7, 2012.

2000 *Mathematics Subject Classification.* Primary 57M50; Secondary 53D10,53D40.

Key words and phrases. contact structure, Reeb dynamics, embedded contact homology, Heegaard Floer homology.

VC supported by the Institut Universitaire de France, ANR Symplexe, ANR Floer Power, and ERC Geodycon. PG supported by ANR Floer Power and ANR TCGD. KH supported by NSF Grants DMS-0805352 and DMS-1105432.

usual, embedded contact homology will be abbreviated as ECH. In [CGH1], we introduced the ECH chain group $\widehat{ECC}(N,\partial N)$ and showed that $\widehat{ECH}(N,\partial N)\simeq\widehat{ECH}(M)$. In the papers [CGH2, CGH3], we defined a chain map

$$\Phi:\widehat{CF}(-M)\to\widehat{ECC}(N,\partial N),$$

which induced an isomorphism

$$\Phi_*:\widehat{HF}(-M)\stackrel{\sim}{\to}\widehat{ECH}(M).$$

The goal of this paper is to extend the above result and prove the following theorem:

Theorem 1.0.1. *If M is a closed oriented 3-manifold, then there is a chain map*

$$\Phi^+:CF^+(-M)\stackrel{\sim}{\to}ECC(M)$$

which is a quasi-isomorphism and which commutes with the U-maps up to homotopy.

We use $\mathbb{F}=\mathbb{Z}/2\mathbb{Z}$ coefficients for both Heegaard Floer homology and ECH.

Remark 1.0.2. The construction of Φ^+ can be carried out with twisted coefficients as in Sections I.6.4 and I.7.1 of [CGH2].

Let $(S,\mathfrak{h})$ be an open book decomposition for M, where S is a genus $g\geq 2$ bordered surface with connected boundary and $\mathfrak{h}\in\mathrm{Diff}(S,\partial S)$.[1] In particular we identify

$$M\simeq(S\times[0,1])/\sim,$$

where $(x,1)\sim(\mathfrak{h}(x),0)$ for all $x\in S$ and $(x,t)\sim(x,t')$ for all $x\in\partial S$ and $t,t'\in[0,1]$. We write $S_t=S\times\{t\}$ for $t\in[0,1]$. Let $\Sigma=S_0\cup -S_{1/2}$ be the Heegaard surface corresponding to $(S,\mathfrak{h})$.

Given a pair $(\Sigma_0,\mathfrak{h}_0)$ consisting of a surface Σ_0 and $\mathfrak{h}_0\in\mathrm{Diff}(\Sigma_0)$, we write the mapping torus of $(\Sigma_0,\mathfrak{h}_0)$ as:

$$N_{(\Sigma_0,\mathfrak{h}_0)}=(\Sigma_0\times[0,2])/(x,2)\sim(\mathfrak{h}_0(x),0).$$

The map Φ, defined in Section I.6.2, is induced by the cobordism W_+ which is an S_0-fibration and which restricts to a half-cylinder over $[0,1]\times S_0$ at the positive end and to a half-cylinder over the suspension $N_{(S_0,\mathfrak{h})}$ at the negative end. We say that W_+ is a cobordism "from $[0,1]\times S_0$[2] to $N_{(S_0,\mathfrak{h})}$."

The map Φ^+ is induced by a cobordism W^+ from $[0,1]\times\Sigma$ to M which extends W_+ and is described below.[3] Although Φ was defined in terms of just one page S_0, we can no longer ignore the $S_{1/2}$ portion of Σ when defining Φ^+, since we do not know how to express $HF^+(-M)$ in terms of S_0.

---

[1] The condition $g\geq 2$ is a technical condition which will used in the definition of Φ^+.

[2] We will interchangeably write $[0,1]\times S_0$ and $S_0\times[0,1]$. This is partly due to the fact that the open book is usually written as $(S\times[0,1])/\sim$ and the positive end of W_+ is a "symplectization" $\mathbb{R}\times[0,1]\times S_0$.

[3] The reader is warned that we are distinguishing W_+ from W^+, i.e., between lower and upper indices.

A symplectic cobordism similar to W^+ is constructed by Wendl in [We4].

1.0.1. *The cobordism W^+.* We give a description of $W^+ = W_0^+ \cup W_1^+ \cup W_2^+$ and $W_+ = W_{+,0} \cup W_{+,1} \cup W_{+,2}$ as topological spaces, where $W_{+,i} \subset W_i^+$ for $i = 0, 1, 2$. See Figure 1. *The description given here is the simplified version of the actual construction, and the notation of Section 1.0.1 is not used outside of Section 1.0.1.*



FIGURE 1. Schematic diagram for $W_0^+ \cup W_1^+$ which indicates the fibers over each subsurface.

First extend $\mathpzc{h} \in \operatorname{Diff}(S_0, \partial S_0)$ to $\mathpzc{h}^+ \in \operatorname{Diff}(\Sigma)$ so that $\mathpzc{h}^+|_{S_{1/2}} = id$. Let $N_{(\Sigma, \mathpzc{h}^+)}$ and $N_{(S_0, \mathpzc{h})}$ be the mapping tori of $\mathpzc{h}^+$ and $\mathpzc{h}$ and let

$$\pi : [0,\infty) \times N_{(\Sigma,\mathpzc{h}^+)} \to [0,\infty) \times \mathbb{R}/2\mathbb{Z}$$

be the projection $(s, x, t) \mapsto (s, t)$. Then define $B_+^0 = ([0,\infty) \times \mathbb{R}/2\mathbb{Z}) - B_+^c$, where B_+^c is the subset $[2,\infty) \times [1, 2]$ with the corners rounded. We then set

$$W_0^+ := \pi^{-1}(B_+^0), \quad W_{+,0} := \pi^{-1}(B_+^0) \cap ([0,\infty) \times N_{(S_0,\mathpzc{h})}).$$

Next we set

$$W_1^+ := S_{1/2} \times D^2, \quad W_{+,1} := \varnothing$$

and identify $\{0\} \times S_{1/2} \times \mathbb{R}/2\mathbb{Z} \subset \partial W_0^+$ with $S_{1/2} \times \partial D^2 \subset \partial W_1^+$ via the map $(0, x, t) \mapsto (x, e^{\pi i t})$. Then one component of $\partial(W_0^+ \cup W_1^+)$ is given by $M = (\{0\} \times N_{(S_0,\mathpzc{h})}) \cup (\partial S_0 \times D^2)$.

Finally we set

$$W_2^+ := (-\infty, 0] \times M, \quad W_{+,2} := (-\infty, 0] \times (\{0\} \times N_{(S_0,\mathpzc{h})}),$$

where $\{0\} \times M$ is identified with M.

1.0.2. *Sketch of proof.* The proof of Theorem 1.0.1 proceeds as follows:

Step 1. Express the U-map on $HF^+(-M)$ as a count of $I_{HF} = 2$ curves that pass through a point, in analogy with the definition of U in ECH. This is given by Theorem 3.1.4.

Step 2. Construct a symplectic cobordism (W^+,Ω^+) from $[0,1]\times\Sigma$ to M, together with stable Hamiltonian and contact structures on $[0,1]\times\Sigma$ and M. This is the goal of Section 4.

Step 3. Define the chain map Φ^+ as a count of $I_{W^+}=0$ curves in W^+ and show that Φ^+ commutes with the U-maps on both sides up to a chain homotopy K. This is done in Section 5.

Step 4. By an algebraic theorem (Theorem 6.1.5), Φ^+ is a quasi-isomorphism if a map

$$\Phi_{alg}:\widehat{CF}(-M)\to\widehat{ECC}(M),$$

defined using Φ^+ and K, is a quasi-isomorphism.

Step 5. By Theorem 6.4.1, the map Φ_{alg} is a quasi-isomorphism. This is proved by relating Φ_{alg} to the quasi-isomorphism Φ from [CGH2, CGH3].

2. HEEGAARD FLOER CHAIN COMPLEXES

The goal of this section is to introduce some notation and recall the definition of the chain complex $CF^+(\Sigma,\boldsymbol{\alpha},\boldsymbol{\beta},z^f,J)$, whose homology is $HF^+(-M)$.

2.1. **Heegaard data.**

Let M be a closed oriented 3-manifold and let (S,h) be an open book decomposition for M.

We use the following notation, which is similar to that of Section I.4.9.1:

- $\Sigma=S_0\cup -S_{1/2}$ is the associated genus $2g$ Heegaard surface of M;
- $\mathbf{a}=\{a_1,\dots,a_{2g}\}$ is a basis of arcs for S and $\mathbf{b}$ is a small pushoff of $\mathbf{a}$ as given in Figure I.1;
- x_i and x_i' are the endpoints of a_i in ∂S_0 that correspond to the coordinates of the contact class and x_i'' is the unique point of $a_i\cap b_i\cap int(S_{1/2})$;
- $\boldsymbol{\alpha}=(\mathbf{a}\times\{\frac{1}{2}\})\cup(\mathbf{a}\times\{0\})$ and $\boldsymbol{\beta}=(\mathbf{b}\times\{\frac{1}{2}\})\cup(h(\mathbf{a})\times\{0\})$ are the collections of compressing curves on the Heegaard surface Σ;
- z^f is a point in the large (i.e., non-thin-strip) component of $S_{1/2}-\boldsymbol{\alpha}-\boldsymbol{\beta}$ and $(z')^f$ is a point which is close but not equal to z^f.

We say that the pointed Heegaard diagram $(\Sigma,\boldsymbol{\alpha},\boldsymbol{\beta},z^f)$ is *compatible with* (S,h).

Remark 2.1.1. *The orientation for Σ is opposite to that of Section I.4.9.1.* This is done so that the triple $(S,\mathbf{a},h(\mathbf{a}))$, used in [CGH2, CGH3], embeds in $(\Sigma,\boldsymbol{\alpha},\boldsymbol{\beta})$ in an orientation-preserving manner.

2.2. **Symplectic data.**

The stable Hamiltonian structure on $[0,1]\times\Sigma$ with coordinates (t,x) is given by (λ,ω), where $\lambda=dt$ and ω is an area form on Σ which makes $(\boldsymbol{\alpha},\boldsymbol{\beta},z^f)$ *weakly admissible with respect to* ω, i.e., each periodic domain has zero ω-area. The plane field $\xi=\ker\lambda$ is equal to the tangent plane field of $\{t\}\times\Sigma$ and the Hamiltonian vector field is $R=\frac{\partial}{\partial t}$.

We introduce the "symplectization"

$$(W,\Omega)=(\mathbb{R}\times[0,1]\times\Sigma, ds\wedge dt+\omega),$$

where (s,t) are coordinates on $\mathbb{R}\times[0,1]$. Let $\pi_B : W \to B = \mathbb{R}\times[0,1]$ be the projection along the fibers $\{(s,t)\}\times\Sigma$.

Let J be an Ω-admissible almost complex structure on W; we assume that J is regular (cf. Lemma I.4.7.2 and [Li, Proposition 3.8]). We also define the Lagrangian submanifolds

$$L_{\boldsymbol{\alpha}} = \mathbb{R}\times\{1\}\times\boldsymbol{\alpha}, \quad L_{\boldsymbol{\beta}} = \mathbb{R}\times\{0\}\times\boldsymbol{\beta}.$$

2.3. **The chain complex $CF^+(\Sigma,\boldsymbol{\alpha},\boldsymbol{\beta},z^f,J)$.** In this subsection we recall the definition of the chain complex $CF^+(\Sigma,\boldsymbol{\alpha},\boldsymbol{\beta},z^f,J)$, whose homology group

$$HF^+(\Sigma,\boldsymbol{\alpha},\boldsymbol{\beta},z^f,J)$$

is isomorphic to $HF^+(-M)$. This definition is due to Lipshitz [Li], with one modification: we are using the ECH index I_{HF} from Definition I.4.5.11. We will often suppress J from the notation.

Let $\mathcal{S}=\mathcal{S}_{\boldsymbol{\alpha},\boldsymbol{\beta}}$ be the set of $2g$-tuples $\mathbf{y}=\{y_1,\dots,y_{2g}\}$ of intersection points of $\boldsymbol{\alpha}$ and $\boldsymbol{\beta}$ for which there exists some permutation $\sigma\in\mathfrak{S}_{2g}$ such that $y_j\in\alpha_j\cap\beta_{\sigma(j)}$ for all j. Then $CF^+(\Sigma,\boldsymbol{\alpha},\boldsymbol{\beta},z^f,J)$ is generated over $\mathbb{F}$ by pairs $[\mathbf{y},i]$, where $\mathbf{y}\in\mathcal{S}$ and $i\in\mathbb{N}$.

The differential $\partial=\partial_{HF}$ is given by

$$\partial[\mathbf{y},i] = \sum_{[\mathbf{y}',j]\in\mathcal{S}\times\mathbb{N}} \langle\partial[\mathbf{y},i],[\mathbf{y}',j]\rangle\cdot[\mathbf{y}',j],$$

where the coefficient $\langle\partial[\mathbf{y},i],[\mathbf{y}',j]\rangle$ is the count of index $I_{HF}=1$ finite energy holomorphic multisections in (W,J) with Lagrangian boundary $L_{\boldsymbol{\alpha}}\cup L_{\boldsymbol{\beta}}$ from $\mathbf{y}$ to $\mathbf{y}'$, whose algebraic intersection with the holomorphic strip $\mathbb{R}\times[0,1]\times\{(z')^f\}$ is $(i-j)$. We will often refer to such curves as curves *from* $[\mathbf{y},i]$ *to* $[\mathbf{y}',j]$.

Let us write $\partial=\sum_{k=0}^{\infty}\partial_k$, where ∂_k only counts curves whose algebraic intersection with $\mathbb{R}\times[0,1]\times\{(z')^f\}$ is k.

3. THE GEOMETRIC U-MAP

3.1. **Introduction.** In [OSz1, Li], the U-map

$$U : CF^+(\Sigma,\boldsymbol{\alpha},\boldsymbol{\beta},z^f)\to CF^+(\Sigma,\boldsymbol{\alpha},\boldsymbol{\beta},z^f),$$

is defined algebraically as $U([\mathbf{y},i])=[\mathbf{y},i-1]$. The goal of this section is to give a geometric definition of the U-map which is analogous to that of ECH.

Let $z^f,(z')^f$ be as before and let $z=(z^b,z^f)\in W=B\times\Sigma$, where $z^b\in int(B)$. Let $J^\lozenge$ be a generic C^l-small perturbation of J such that $J^\lozenge=J$ away from a small neighborhood $N(z)\subset W$ of z and such that $N(z)\cap(\mathbb{R}\times[0,1]\times\{(z')^f\})=\varnothing$. In particular, we assume that there are no $J^\lozenge$-holomorphic curves that are homologous to $\{pt\}\times\Sigma$ and pass through z.

Remark 3.1.1. When we refer to "C^l-close" almost complex structures, etc., we assume that $l>0$ is sufficiently large.

Definition 3.1.2 (Geometric U-map)**.** The *geometric U-map with respect to the point z* is the map:

$$U_z([\mathbf{y},i]) = \sum_{[\mathbf{y}',j]\in\mathcal{S}\times\mathbb{N}} \langle U_z([\mathbf{y},i]),[\mathbf{y}',j]\rangle\cdot[\mathbf{y}',j],$$

where the coefficient $\langle U_z([\mathbf{y},i]),[\mathbf{y}',j]\rangle$ is the count of index $I_{HF}=2$ finite energy holomorphic curves in $(W,J^\Diamond)$ with Lagrangian boundary $L_{\boldsymbol{\alpha}}\cup L_{\boldsymbol{\beta}}$ from $[\mathbf{y},i]$ to $[\mathbf{y}',j]$ that pass through z.

By our choice of $J^\Diamond$, "passing through z" is a generic codimension 2 condition, i.e., if u is a simple curve from $[\mathbf{y},i]$ to $[\mathbf{y}',j]$ that passes through z, then $I(u)\geq 2$. Moreover, by a simple ECH index count, an $I(u)\leq 3$ curve that passes through z cannot have a fiber component.

Proposition 3.1.3. *U_z is a chain map.*

Proof. This follows from standard arguments in symplectic geometry by using analytical results proved in [Li]. ☐

Theorem 3.1.4. *There exists a chain homotopy*

$$H: CF^+(\Sigma,\boldsymbol{\alpha},\boldsymbol{\beta},z^f)\to CF^+(\Sigma,\boldsymbol{\alpha},\boldsymbol{\beta},z^f)$$

such that

$$U_z - U = H\circ\partial_{HF}+\partial_{HF}\circ H. \tag{3.1.1}$$

Moreover, for all $\mathbf{y}\in\mathcal{S}$, one has $H([\mathbf{y},0])=0$.

The rest of this section is devoted to the proof of Theorem 3.1.4.

3.2. **A model calculation.** Let Σ be a closed surface of genus k. We consider the manifold $M=D\times\Sigma$, where $D=\{|z|\leq 1\}\subset\mathbb{C}$. Let $\pi_D: M\to D$ and $\pi_\Sigma: M\to\Sigma$ be the projections of M onto the first and second factors. Let $\boldsymbol{\beta}=\{\beta_1,\dots,\beta_k\}$ be the set of β-curves for Σ. Choose $z^f\in\Sigma-\boldsymbol{\beta}$ and let $z=(0,z^f)\in M$.

Let $J=j_D\times j_\Sigma$ be a product complex structure on M and $J^\Diamond$ be a generic C^l-small perturbation of J such that $J^\Diamond=J$ away from a small neighborhood of z. The key feature of $J^\Diamond$ is that all the $J^\Diamond$-holomorphic curves that pass through z are regular.

We then define the moduli space $\mathcal{M}_A(M,J^*)$, $*=\varnothing$ or $\Diamond$, of stable maps

$$u:(F,j)\to(M,J^*)$$

in the class $A=[\{pt\}\times\Sigma]+k[D\times\{pt\}]\in H_2(M,\partial D\times\boldsymbol{\beta})$, such that ∂F has k connected components and each component of ∂F maps to a distinct Lagrangian $\partial D\times\beta_i$, $i=1,\dots,k$. We choose points $w_i\in\beta_i$, $i=1,\dots,k$, and define

$$\mathbf{w}=\{(1,w_1),\dots,(1,w_k)\}\subset M,$$

Then let

$$\mathcal{M}_A(M,J^*;z,\mathbf{w})\subset\mathcal{M}_A(M,J^*)$$

be the subset of curves that pass through z and $\mathbf{w}$. We use the modifier "irr" to denote the subset of irreducible curves.

3.2.1. *ECH index.* We briefly indicate the definition of the ECH index I of a homology class $B \in H_2(M, \partial D \times \boldsymbol{\beta})$ which admits a representative F such that each component of ∂F maps to a distinct $\partial D \times \beta_i$. Although we call I the "ECH index", what we are defining here is a relative version of Taubes' index from [T4].

Let τ be a trivialization of $T\Sigma$ along $\boldsymbol{\beta}$, given by a nonsingular tangent vector field X_1 along $\boldsymbol{\beta}$, and let τ' be a trivialization of TD along ∂D, given by an outward-pointing radial vector field X_2 along ∂D. Let $Q_{(\tau,\tau')}(B)$ be the intersection number between an embedded representative u of B and its pushoff, where the boundary of u is pushed off in the direction given by $J(X_1)$.

Definition 3.2.1. The *ECH index of the homology class* B is:

$$I(B) = c_1(TM|_B, (\tau, \tau')) + \mu_{(\tau,\tau')}(\partial B) + Q_{(\tau,\tau')}(B).$$

The following is the relative version of the adjunction inequality:

Lemma 3.2.2 (Index inequality)**.** *Let* $u : (F, j) \to (M, J^*)$ *be a holomorphic curve in the class* $B \in H_2(M, \partial D \times \boldsymbol{\beta})$*. Then*

$$\operatorname{ind}(u) + 2\delta(u) = I(B),$$

where $\delta(u) \geq 0$ *is an integer count of the singularities.*

Proof. Similar to the proof of Theorem I.4.5.13. □

We now calculate some ECH and Fredholm indices:

Lemma 3.2.3. *If* $B = [\{pt\} \times \Sigma] + k_0[D \times \{pt\}]$ *with* $k_0 \leq k$*, then*

$$I(B) = 2 - 2k + 3k_0.$$

Proof. We compute that

$$\begin{aligned} I(B) &= I([\{pt\} \times \Sigma] + k_0[D \times \{pt\}]) \\ &= I([\{pt\} \times \Sigma]) + k_0 \cdot I([D \times \{pt\}]) + 2k_0 \cdot \langle [\{pt\} \times \Sigma], [D \times \{pt\}] \rangle \\ &= (2 - 2k) + k_0 \cdot 1 + 2k_0 = 2 - 2k + 3k_0. \end{aligned}$$

Here $\langle , \rangle$ denotes the algebraic intersection number. □

Lemma 3.2.4. *If* $B = [\{pt\} \times \Sigma] + k_0[D \times \{pt\}]$ *with* $k_0 \leq k$ *and* u *is an irreducible* $J^\Diamond$*-holomorphic curve in the class* B*, then*

$$\operatorname{ind}(u) = 2 - 2k + 3k_0 - \delta(u).$$

Proof. Follows from Lemma 3.2.3 and the index inequality. □

3.2.2. *Main result.* The following is the main result of this subsection:

Theorem 3.2.5. *If* $J^\Diamond$ *is generic, then the following hold:*

1. $\mathcal{M}_A(M, J^\Diamond; z, \mathbf{w}) = \mathcal{M}_A^{irr}(M, J^\Diamond; z, \mathbf{w})$;
2. $\mathcal{M}_A(M, J^\Diamond; z, \mathbf{w})$ *is compact, regular, and* 0*-dimensional;*
3. *the curves of* $\mathcal{M}_A(M, J^\Diamond; z, \mathbf{w})$ *are embedded; and*
4. $\#\mathcal{M}_A(M, J^\Diamond; z, \mathbf{w}) \equiv 1$ *mod* 2.

Hence $\#\mathcal{M}_A(M, J^\Diamond; z, \mathbf{w})$ is a certain relative Gromov-Witten invariant [IP1] which is computed to be 1 mod 2. (What we are really computing here is a relative Gromov-Taubes invariant [T4], although the two invariants coincide in this case.)

Proof. (1) Let us write $\mathcal{M} = \mathcal{M}_A(M, J^\Diamond; z, \mathbf{w})$. Arguing by contradiction, suppose $u \in \mathcal{M} - \mathcal{M}^{irr}$. Then u consists of an irreducible component u_0 which passes through z and $k_0 < k$ points of $\mathbf{w}$, together with $k - k_0$ copies of $D \times \{pt\}$. By Lemma 3.2.4, $\operatorname{ind}(u_0) \leq 2 - 2k + 3k_0$. On the other hand, the point constraints are $(k_0 + 2)$-dimensional. Hence u_0 does not exist for generic $J^\Diamond$, which is a contradiction.

(2),(3) The compactness follows from the usual Gromov compactness theorem. The regularity of $\mathcal{M}$ is immediate from the genericity of $J^\Diamond$ and (1). Lemma 3.2.4 implies the dimension calculation, as well as (3).

(4) We degenerate the fiber Σ into a union of k tori which are successively attached to one another. We perform this pinching away from the curves $\boldsymbol{\beta}$ and make sure that each torus contains exactly one component of $\boldsymbol{\beta}$. Then by a degeneration/gluing argument as in Section II.2.4.4, it suffices to prove the proposition for $k = 1$. The case $k = 1$ is proved in Lemmas 3.2.8 and 3.2.9. □

3.2.3. *Doubling.* We now explain how to double $u \in \mathcal{M}_A(M, J^\Diamond; z, \mathbf{w})$. For technical reasons we will assume that (Σ, j_Σ) admits an anti-holomorphic involution σ_Σ so that the curve $\boldsymbol{\beta}$ is contained in the fixed point set of σ_Σ.

Let $\mathcal{D}(M) = M_1 \cup M_2$ be the double of $M = D \times \Sigma$, obtained by gluing two copies M_1 and M_2 of M along their boundaries $\partial D \times \Sigma$ via the identification $(x, y)_1 \sim (x, \sigma_\Sigma(y))_2$, where the subscript $i = 1, 2$ indicates the ith copy. Let $\mathcal{S}$ be the involution of $\mathcal{D}(M)$ given by

$$\mathcal{S} : (x, y)_1 \mapsto (x, \sigma_\Sigma(y))_2, \quad \mathcal{S} : (x, y)_2 \mapsto (x, \sigma_\Sigma(y))_1.$$

We then define the almost complex structure $\mathcal{D}(J^\Diamond)$ on $\mathcal{D}(M)$ by taking $J^\Diamond$ on M_1 and $\mathcal{S}(-J^\Diamond)$ on M_2.

Given $u \in \mathcal{M}_A(M, J^\Diamond; z, \mathbf{w})$, let $\mathcal{D}(u)$ be its double, obtained by gluing u and $\mathcal{S}(u)$. The map $\mathcal{D}(u)$ is holomorphic by the Schwarz reflection principle. Therefore it is an element of

$$\mathcal{M}^{irr}_{\mathcal{D}, J^\Diamond} := \mathcal{M}^{irr}_{\mathcal{D}(A)}(\mathcal{D}(M), \mathcal{D}(J^\Diamond); z, \mathcal{S}(z), \mathbf{w}),$$

because it represents the class

$$\mathcal{D}(A) := 2[\{pt\} \times \Sigma] + [S^2 \times \{pt\}]$$

and passes through 3 points: $(1, w)$, z, and $\mathcal{S}(z)$, where $z \in M_1$. Conversely, if $v : (F, j) \to \mathcal{D}(M)$ is in $\mathcal{M}_{\mathcal{D}, J^\Diamond}$, then $\mathcal{S} \circ v : (F, -j) \to \mathcal{D}(M)$ is also in $\mathcal{M}_{\mathcal{D}, J^\Diamond}$. Thus all the curves of $\mathcal{M}_{\mathcal{D}, J^\Diamond}$ come in pairs, except those that are $\mathcal{S}$-invariant, and the $\mathcal{S}$-invariant curves are those obtained by the doubling procedure.

Summarizing, we have:

Lemma 3.2.6. $\#\mathcal{M}^{irr}_A(M, J^\Diamond; z, \mathbf{w}) \equiv \#\mathcal{M}^{irr}_{\mathcal{D}, J^\Diamond}$ *mod* 2.

3.2.4. *The $k = 1$ case.* For the rest of the subsection we assume $k = 1$. We first consider the case where $J = j_D \times j_\Sigma$ is a product complex structure.

Lemma 3.2.7. *If $k = 1$, then:*

1. $\mathcal{M}_A(M, J; z, \mathbf{w})$ *is a one-element set consisting of a degenerate curve* $(D \times \{w\}) \cup (\{0\} \times \Sigma)$*; and*
2. $\mathcal{M}_{\mathcal{D},J}$ *is a one-element set consisting of a degenerate curve* $(S^2 \times \{w\}) \cup (\{0\} \times \Sigma)_1 \cup (\{0\} \times \Sigma)_2$.

Proof. (1) follows from the homological constraint

$$A = [\{pt\} \times \Sigma] + [D \times \{pt\}].$$

If $u : (F, j) \to (M, J)$ is a stable map in $\mathcal{M}_A(M, J; z, \mathbf{w})$, then $\pi_D \circ u$ and $\pi_\Sigma \circ u$ are degree 1 maps. This implies that F consists of two components F_1, F_2 and $\pi_D \circ u|_{F_1}$ and $\pi_\Sigma \circ u|_{F_2}$ are biholomorphisms. On the other hand, $\pi_\Sigma \circ u|_{F_1}$ maps to a point since F_1 is a disk and $\pi_D \circ u|_{F_2}$ maps to a point since otherwise the cardinality of $(\pi_D \circ u)^{-1}(pt)$ for generic pt will be larger than $\deg(\pi_D \circ u) = 1$. (2) is similar. ☐

Let $J^\diamondsuit$ be an almost complex structure which is C^l-close to J. By Gromov compactness and Lemma 3.2.7, all the curves of $\mathcal{M}_A(M, J^\diamondsuit; z, \mathbf{w})$ and $\mathcal{M}_{\mathcal{D},J^\diamondsuit}$ are close to the degenerate curves described in Lemma 3.2.7.

Lemma 3.2.8. *If $k = 1$ and $J^\diamondsuit$, $\mathbf{w}$, and $\boldsymbol{\beta}$ are generic, then the following hold:*

1. $\mathcal{M}_{\mathcal{D},J^\diamondsuit} = \mathcal{M}^{irr}_{\mathcal{D},J^\diamondsuit}$;
2. *the curves of* $\mathcal{M}_{\mathcal{D},J^\diamondsuit}$ *are embedded; and*
3. $\mathcal{M}_{\mathcal{D},J^\diamondsuit}$ *is compact, regular, and* 0*-dimensional.*

Proof. (1) Arguing by contradiction, suppose $v \in \mathcal{M}_{\mathcal{D},J^\diamondsuit} - \mathcal{M}^{irr}_{\mathcal{D},J^\diamondsuit}$. If v has a component $S^2 \times \{pt\}$, then v must also have a fiber component $\widetilde{v}$ which is close to $(\{0\} \times \Sigma)_i$, $i = 1$ or 2, since v is close to a degenerate curve from Lemma 3.2.7(2). On the other hand, if v does not have a component $S^2 \times \{pt\}$, then v must have a fiber component $\widetilde{v}$ which is close to some $(\{0\} \times \Sigma)_i$. In either case, such a fiber component $\widetilde{v}$ cannot exist by the genericity of $J^\diamondsuit$. This proves (1).

(2) Let $v \in \mathcal{M}^{irr}_{\mathcal{D},J^\diamondsuit}$. The proof is similar to that of Lemma 3.2.5(3) and follows from the adjunction inequality [M1, M2] (compare with Lemma 3.2.2): If

$$I(v) = c_1(v^*T\mathcal{D}(M))) + Q(v),$$

where $Q(v)$ is the self-intersection number of v, then

$$\operatorname{ind}(v) + 2\delta(v) = I(v),$$

where $\delta(v) \geq 0$ is an integer count of the singularities. Since $c_1(v^*T\mathcal{D}(M))) = 2$ and $Q(v) = 4$, it follows that $I(v) = 6$. On the other hand,

$$\operatorname{ind}(v) = -\chi(F) + 2c_1(v^*T\mathcal{D}(M)) = -(-2) + 2(2) = 6,$$

where F is the domain of v with $\chi(F) = -2$. Hence v is embedded by the adjunction inequality.

(3) Since v is embedded by (2) and $c_1(v^*T\mathcal{D}(M))) = 2$, the regularity of v without the point constraints follows from automatic transversality (cf. Hofer-Lizan-Sikorav [HLS, Theorem 1]). The regularity with point constraints is the consequence of the genericity of $J^\Diamond$, $\mathbf{w}$, and $\boldsymbol{\beta}$. The rest of the assertion is immediate. ☐

Lemma 3.2.9. *If $k = 1$ and $J^\Diamond$, $\mathbf{w}$, and $\boldsymbol{\beta}$ are generic, then $\#\mathcal{M}_A(M, J^\Diamond; z, \mathbf{w}) \equiv 1$ mod 2.*

Proof. By Lemma 3.2.6, Lemma 3.2.8(1), and Theorem 3.2.5(1),

$$\#\mathcal{M}_A(M, J^\Diamond; z, \mathbf{w}) \equiv \#\mathcal{M}_{\mathcal{D},J^\Diamond} \text{ mod } 2. \tag{3.2.1}$$

We reduce the calculation of $\mathcal{M}_{\mathcal{D},J^\Diamond}$ to a calculation in McDuff-Salamon [MS, Example 8.6.12] by degenerating the base S^2 into two spheres along the curve corresponding to the boundary of the two copies of D.

More precisely, let

$$B = [S^2 \times \{pt\}] + [\{pt\} \times \Sigma] \in H_2(S^2 \times \Sigma),$$

let I be a product complex structure on $S^2 \times \Sigma$, and let $I^\Diamond$ be a C^l-small perturbation of I such that $I = I^\Diamond$ away from a neighborhood of $(0, z^f)$. Here we are viewing $S^2 \simeq \mathbb{C} \cup \{\infty\}$. Let

$$\mathcal{M}_B := \mathcal{M}_B(S^2 \times \Sigma, I^\Diamond; (0, z^f), (\infty, w))$$

be the moduli spaces of $I^\Diamond$-curves in the class B that pass through $(0, z^f)$ and (∞, w), and let $\mathcal{M}'_B$ be the subset of curves in $\mathcal{M}_B$ that are contained in a neighborhood of $(S^2 \times \{w\}) \cup (\{0\} \times \Sigma)$. By McDuff-Salamon [MS, Example 8.6.12], $\#\mathcal{M}'_B = 1$. By the symplectic sum theorem of Ionel-Parker [IP2, p. 940] (also see Section II.2.4.4),

$$\#\mathcal{M}_{\mathcal{D},J^\Diamond} \equiv (\#\mathcal{M}'_B)^2 \equiv 1 \text{ mod } 2.$$

Hence $\#\mathcal{M}_A(M, J^\Diamond; z, \mathbf{w}) \equiv 1$ mod 2 by Equation (3.2.1). ☐

3.3. **Family of cobordisms.** We now describe a family of marked points $z_\tau \in W$ and a family of almost complex structures $J^\Diamond_\tau$ on W for $\tau \in [0, 1)$, as well as their limits for $\tau = 1$. These families give rise to the chain homotopy H of Theorem 3.1.4.

Let $z^b_\tau \in int(B)$, $\tau \in [0, 1)$, be a family of points such that $z^b_0 = z^b$, $\lim_{\tau\to 1} z^b_\tau = (0, 0)$, and $z^b_\tau \in \{s = 0\}$ for $\tau \in [\frac{1}{2}, 1)$. Then let $z_\tau = (z^b_\tau, z^f) \in W$.

Assume that the almost complex structure J on W is a product complex structure on $\mathbb{R} \times [0, \varepsilon] \times \Sigma$ for $\varepsilon > 0$ small. We then define a family of C^l-small perturbations $J^\Diamond_\tau$, $\tau \in [0, 1)$, of J such that $J^\Diamond_\tau = J$ away from a small neighborhood $N(z_\tau)$ of z_τ and

$$N(z_\tau) \cap (\mathbb{R} \times [0, 1] \times \{(z')^f\}) = \varnothing.$$

In the limit $\tau = 1$, the base B_1 is $(B \sqcup D)/\sim$, where $D = \{|z| \leq 1\} \subset \mathbb{C}$ and $\sim$ identifies $(0, 0) \in B$ with $-1 \in D$, and the total space W^1 is $(W \sqcup (D \times \Sigma))/\sim$, where $((0, 0), x) \sim (-1, x)$ for all $x \in \Sigma$. See Figure 2. We write w^b for the node

$[(0,0)] = [-1] \in B_1$. Let $\pi_B : W \to B$ and $\pi_D : D \times \Sigma \to D$ be the projections onto the first factors.



FIGURE 2

The limit z_1 of z_τ is in $D \times \Sigma$ and we assume that $z_1^b = 0 \in int(D)$. When $\tau = 1$, the almost complex structure $J_1^\lozenge$ restricts to the complex structure J on W and to the almost complex structure $(J^{1,D})^\lozenge$, where $J^{1,D}$ is a product complex structure on $D\times\Sigma$ and $(J^{1,D})^\lozenge$ is a C^l-small perturbation of $J^{1,D}$ such that $(J^{1,D})^\lozenge = J^{1,D}$ away from a small neighborhood $N(z_1)$ of z_1 and

$$N(z_1) \cap (D \times \{(z')^f\}) = \varnothing.$$

The Lagrangian boundary condition for $\tau \in [0,1)$ is $L_{\boldsymbol\alpha} \cup L_{\boldsymbol\beta}$. In the limit $\tau = 1$, we use $L_{\boldsymbol\alpha} \cup L_{\boldsymbol\beta}$ for W and $\partial D \times \boldsymbol\beta$ for $D \times \Sigma$.

3.4. **Proof of Theorem 3.1.4.** Let u_{τ_i}, $\tau_i \to 1$, be a sequence of $I_{HF} = 2$ curves in $(W^{\tau_i}, (J^{\tau_i})^\lozenge)$ from $[\mathbf{y}, i]$ to $[\mathbf{y}', i-k]$ that pass through z_{τ_i}. Applying Gromov compactness, we obtain the limit $u_1 = u_B \cup u_D$, where $u_B \subset W$, $u_D \subset D \times \Sigma$, and u_D passes through z_1. Components of u_1 that map to the fiber $\{w^b\} \times \Sigma$ will be viewed as components of u_D.

Lemma 3.4.1.

1. $[u_D] = k_0[\{pt\} \times \Sigma] + 2g[D \times \{pt\}] \in H_2(D \times \Sigma)$ *for some* $0 < k_0 \le k$.
2. $I(u_D) = 2k_0 + 2g \ge 2g + 2$.

Proof. (1) $\deg(\pi_D \circ u_D) = 2g$, since u_{τ_i} is a degree $2g$ multisection of W for each τ_i, away from a neighborhood of $z^b_{\tau_i}$. Also, since $\langle u_{\tau_i}, B \times \{(z^f)'\}\rangle = k$ for all τ_i, it follows that $\langle u_D, D \times \{(z^f)'\}\rangle = k_0$, where $0 < k_0 \le k$. Here $k_0 > 0$ since u_D passes through z_1.

(2) is a consequence of (1) and computations as in the proof of Lemma 3.2.3. We remind the reader that the genus of Σ is $2g$. □

Lemma 3.4.2. $I(u_D) = 2g + 2$ *and* $I_{HF}(u_B) = 0$. *In particular,* $\mathbf{y} = \mathbf{y}'$, u_B *consists of* $2g$ *trivial strips, and* $k_0 = k = 1$.

Proof. The gluing constraints give $I_{HF}(u_\tau) = I(u_D) + I_{HF}(u_B) - 2g = 2$. By the regularity of J and the index inequality, we have $I_{HF}(u_B) \ge 0$. The first sentence of the lemma then follows from Lemma 3.4.1(2); the second sentence is a consequence of the first. □

The first sentence of Theorem 3.1.4 follows from the usual construction of chain homotopies in Floer theory. By Lemma 3.4.2, U_z is chain homotopic to aU, where a is the count of holomorphic curves u_D in $(D \times \Sigma, (J^{1,D})^\Diamond)$ that pass through z_1 and $\mathbf{w} = ((1,y_1),\dots,(1,y_{2g}))$, where $\mathbf{y} = \{y_1,\dots,y_{2g}\}$. Since $a = 1$ modulo 2 by Theorem 3.2.5, U_z is chain homotopic to U.

Next we prove the second sentence of Theorem 3.1.4. For all $\mathbf{y} \in \mathcal{S}$, $H([\mathbf{y},0])$ is obtained by counting $I_{HF} = 1$ curves that pass through z_τ for some $\tau \in (0,1)$ and that do not cross the holomorphic strip $\mathbb{R}\times[0,1]\times\{(z')^f\}$. There are no such curves since $\mathbb{R}\times[0,1]\times\{z^f\}$ is holomorphic and homologous to $\mathbb{R}\times[0,1]\times\{(z')^f\}$: if a curve passes through z_τ, its intersection with $\mathbb{R}\times[0,1]\times\{z^f\}$ is strictly positive by the positivity of intersections, and so is its intersection with $\mathbb{R}\times[0,1]\times\{(z')^f\}$.

4. THE COBORDISM W^+

In this section we give the construction of the symplectic cobordism (W^+,Ω^+) from $[0,1]\times\Sigma$ to M, together with the Lagrangian submanifold $L_{\boldsymbol{\alpha}} \subset \partial W^+$.

4.1. **Construction of** (W^+,Ω^+).

We describe the construction of W^+, leaving some key details for later[4]: First we construct fibrations $\pi_0 : W_0^+ \to B_+^0$ and $\pi_1 : W_1^+ \to D^2$ with fibers diffeomorphic to Σ and $S_{1/2}$. Here $B_+^0 = ([0,\infty)\times\mathbb{R}/2\mathbb{Z}) - B_+^c$ with coordinates (s,t) and B_+^c is the subset $[2,\infty)\times[1,2]$ with the corners rounded. We then glue W_0^+ and W_1^+ and smooth a boundary component $\mathcal{B}$ of $W_0^+\cup W_1^+$ to obtain $\widetilde{\mathcal{B}} \simeq M$. Finally we attach the negative end $W_2^+ = (-\infty,0]\times\widetilde{\mathcal{B}}$ to obtain W^+.

Let $\delta > 0$ be a small irrational number and N a large positive number which depends on δ and whose dependence will be described later.

Lemma 4.1.1. *There exists a symplectic manifold (W^+,Ω^+) which depends on $\delta > 0$ and which satisfies the following:*

1. *There is a symplectic surface $S_{z^f} := \{z^f\}\times(B_+^0\cup D^2)$, obtained by gluing sections $\{z^f\}\times B_+^0 \subset W_0^+$ and $\{z^f\}\times D^2\subset W_1^+$.*
2. *$\Omega^+ = d\Theta^+$ for some 1-form Θ^+ on $W^+ - N(S_{z^f})$, where $N(S_{z^f})$ is a small neighborhood of S_{z^f}.*
3. *Θ^+ is exact on the Lagrangian submanifold $L_{\boldsymbol{\alpha}} \subset \partial W^+$.*
4. *On the positive end*
$$\pi_0^{-1}([3,\infty)\times[0,1]) = [3,\infty)\times\Sigma\times[0,1] \subset W_0^+$$
of W^+, Ω^+ restricts to $\widetilde{\omega}+ds\wedge dt$, where $\widetilde{\omega}$ is an area form on Σ. Moreover,
$$L_{\boldsymbol{\alpha}} \cap \{s\geq 3\} = ([3,\infty)\times\{0\}\times\boldsymbol{\beta}')\cup([3,\infty)\times\{1\}\times\boldsymbol{\alpha}),$$
where $\boldsymbol{\beta}'$ is isotopic to $\boldsymbol{\beta}$.
5. *On the negative end W_2^+ of W^+, Ω^+ restricts to the negative symplectization of a contact form λ_- on $\widetilde{\mathcal{B}} \simeq M$ which is adapted to the open book decomposition $(S,\hbar)$.*

---

[4] Compare with the description in Section 1.0.1, keeping in mind that the notation will be slightly different.

(6) *The manifold $\widetilde{\mathcal{B}} \simeq M$ admits a decomposition into three disjoint pieces: the suspension $N_{(S_0,\hslash_0)}$, where $\hslash_0$ is isotopic to $\hslash$ relative to ∂S_0, a closed neighborhood $N(K)$ of the binding K, and an open thickened torus $\mathcal{N}$ in between that we refer to as the "no man's land".*

(7) *All the orbits of the Reeb vector field R_{λ_-} of λ_- in $int(N(K)) \cup \mathcal{N}$ have λ_--action $\geq \frac{1}{2\delta} - \kappa$, where $\kappa > 0$ is independent of δ. Moreover, $T_+ = \partial N(K)$ (resp. $T_- = \partial N_{(S_0,\hslash_0)}$) is a positive (resp. negative) Morse-Bott torus of meridian orbits.*

(8) *There is an embedding of W_+, defined in Section I.5.1.1, into W^+ such that the restriction $\pi_1 : W_+ \cap W_0^+ \to B_+^0$ is a fibration with fiber S_0, $W_+ \cap W_1^+ = \varnothing$, $W_+ \cap W_2^+ = [0,\infty) \times N_{(S_0,\hslash_0)}$, and W_+ is disjoint from $N(S_{z^f})$.*

Here W^+, Ω^+, Θ^+, $L_{\boldsymbol{\alpha}}$, and λ_- depend on $\delta > 0$.

The S^1-family $\mathcal{P}_+$ (resp. $\mathcal{P}_-$) of simple orbits of T_+ (resp. T_-) can be viewed equivalently as a pair e', h' (resp. e, h) consisting of an elliptic orbit and a hyperbolic orbit. The proof of Lemma 4.1.1 will be given in Section 4.3.

Let $A_{[-1,N]} \simeq [-1,N] \times S^1$ be a small neighborhood of $\partial S_0 = \{0\} \times S^1$ in Σ with coordinates (r_1,θ_1), such that $z^f \notin A_{[-1,N]}$, $A_{[-1,0]} \subset S_0$ and $A_{[0,N]} \subset S_{1/2}$. Here we write $A_{\mathfrak{I}} = \mathfrak{I} \times S^1$ if $\mathfrak{I}$ is a subset of $[-1,N]$. Also let $N(z^f) \subset S_{1/2} - A_{[0,N]} - \boldsymbol{\alpha} - \boldsymbol{\beta}$ be a small ball $D_\tau = \{r' \leq \tau\}$ about z^f, where we are using polar coordinates (r',θ').

The actual construction of (W^+,Ω^+) is a bit involved, and consists of several steps.

Step 1. The following lemma is a rephrasing of Lemma I.2.1.1 and its proof.

Lemma 4.1.2. *After possibly isotoping $\hslash$ relative to ∂S_0, there exists a factorization $\hslash = \hslash_0 \circ \hslash_1$ and a contact form $\lambda = f_t(x)dt + \beta_t(x)$, $(x,t) \in S_0 \times [0,2]$, on $N_{(S_0,\hslash_0)}$ with Reeb vector field R_λ, such that the following hold:*

(1) *$\hslash : S_0 \times \{0\} \xrightarrow{\sim} S_0 \times \{0\}$ is the first return map of R_λ.*
(2) *$\hslash$ has no elliptic periodic point of period $\leq 2g$ in $int(S_0)$, as required for technical reasons in II.1.0.1.*
(3) *$\hslash_0 = id$ on $A_{[-1/2,0]}$.*
(4) *$\hslash_1$ is the flow of R_λ from $S_0 \times \{0\}$ to $S_0 \times \{2\}$.*
(5) *R_λ is parallel to ∂_t on $(S_0 - A_{[-1,0]}) \times [0,2]$. In particular, $\hslash_1 = id$ on $S_0 - A_{[-1,0]}$.*
(6) *$f_t(r_1,\theta_1) = 1 + \varepsilon r_1^2/2$ and $\beta_t(r_1,\theta_1) = (C + r_1)d\theta_1$ on $A_{[-1/2,0]}$, for $\varepsilon > 0$ sufficiently small and $C > 0$. In particular, f_t and β_t are independent of t and R_λ is parallel to $\partial_t - \varepsilon r_1 \partial_\theta$ on $A_{[-1/2,0]}$.*
(7) *$|d_2 f_t|_{A_{[-1/2,0]}}|_{C^0} \leq \delta$ and $\frac{1}{2} \leq f_t \leq 2$.*

Here $\varepsilon > 0$ depends on $\delta > 0$, d_2 is the differential in the S_0-direction, and the C^0-norm is with respect to a fixed Riemannian metric on S_0.

Step 2. We then extend $\hbar_0, \hbar_1, \hbar \in \mathrm{Diff}(S_0, \partial S_0)$ to $\hbar_0^+, \hbar_1^+, \hbar^+ = \hbar_0^+ \circ \hbar_1^+ \in \mathrm{Diff}(\Sigma)$ and the contact form λ to the contact form $\lambda_+ = f_t dt + \beta_t$ to $N_{(\Sigma - N(z^f), \hbar_0^+)}$, all of which depend on $\delta > 0$, as follows:

- (3') $\hbar_0^+ = id$ on $S_{1/2}$.
- (4') $\hbar_1^+|_{\Sigma - N(z^f)}$ is the flow of R_{λ_+} from $(\Sigma - N(z^f)) \times \{0\}$ to $(\Sigma - N(z^f)) \times \{2\}$ and $\hbar_1^+|_{N(z^f)} = id$.
- (5') f_t and β_t are independent of t on $S_{1/2} - N(z^f)$. Hence R_{λ_+} is parallel to $\partial_t + X_f$, where X_f is the Hamiltonian vector field satisfying $i_{X_f}\omega = d_2 f$ and ω is an area form on Σ which agrees with $d_2\beta_t$ on $\Sigma - N(z^f)$.
- (6a') $f_t(r', \theta') = const > 0$ and $\beta_t(r', \theta') = (-C' + r')d\theta'$ near $\partial N(z^f)$, for $-C' > 0$. In particular, R_{λ_+} is parallel to ∂_t near the suspension of $\partial N(z^f)$.
- (6b') $f_t(r_1, \theta_1) = 1 + \varepsilon r_1^2/2$ near $A_{\{0\}}$ and $\beta_t(r_1, \theta_1) = (C + r_1)d\theta_1$ on $A_{[0,N]}$.
- (7') $|d_2 f_t|_{S_{1/2} - N(z^f)}|_{C^0} \leq \delta$ and $\frac{1}{2} \leq f_t|_{S_{1/2} - N(z^f)} \leq 2$.

Without loss of generality we may assume that $\boldsymbol{\alpha} \times \{1\}$ is Legendrian with respect to λ_+. This is an easy consequence of the Legendrian realization principle; see for example [H, Theorem 3.7].

Step 3 (Construction of (W_0^+, Ω_0^+)). Let

$$\widetilde{W_0^+} = ([0, \infty) \times \Sigma \times [0, 2])/(s, x, 2) \sim (s, \hbar_0^+(x), 0)$$

and let $\pi_0 : \widetilde{W_0^+} \to [0, \infty) \times \mathbb{R}/2\mathbb{Z}$ be the projection $(s, x, t) \mapsto (s, t)$. We then set

$$W_0^+ := \pi_0^{-1}(B_+^0).$$

Let $g : [0, \frac{1}{2}] \to \mathbb{R}$ be a smooth function such that $g(r) = 1 + \varepsilon r^2/2$ near $r = 0$, $0 < g'(r) \leq \delta$ for $r \in (0, \frac{1}{2})$, $g'(\frac{1}{2}) = 0$, and $g(\frac{1}{2}) = 1 + \varepsilon$. In particular, this requires $2\varepsilon < \delta$. Then let

$$\lambda_{+,s} = f_{s,t} dt + \beta_t, \quad s \in [0, \infty),$$

be a 1-parameter family of contact forms[5] on $N_{(\Sigma - N(z^f), \hbar_0^+)}$ such that the following hold:

- (a) $\lambda_{+,s} = \lambda_+$ if $s \geq \frac{3}{2}$ or $(x, t) \in N_{(S_0, \hbar_0)}$.
- (b) $\lambda_{+,s}$ is independent of s if $s \in [0, \frac{1}{2}]$.
- (c) $f_{0,t}(r_1, \theta_1) = g(r_1)$ on $A_{[0,1/2]}$.
- (d) $f_{0,t}|_{S_{1/2} - A_{[0,1/2]} - N(z^f)} = 1 + \varepsilon$. In particular, $d\lambda_{+,0} = d_2\beta_t$ and $R = \partial_t$ on the suspension of $S_{1/2} - A_{[0,1/2]} - N(z^f)$.
- (e) $f_{s,t}$ is a constant $C_s > 0$ near $\partial N(z^f)$.
- (f) $|d_2 f_{s,t}|_{A_{[-1/2,0]} \cup S_{1/2} - N(z^f)}|_{C^0} \leq \delta$, $|\partial_s f_{s,t}|_{A_{[-1/2,0]} \cup S_{1/2} - N(z^f)}|_{C^0} \leq \delta$ and $\frac{1}{2} \leq f_{s,t}|_{\Sigma - N(z^f)} \leq 2$ for all s, t.

---

[5] Note that β_t does not depend on s.

We then define:

$$\Omega_0^+ := \widetilde{\omega} + ds \wedge dt,$$

where

$$\widetilde{\omega} = \begin{cases} d\lambda_{+,s} & \text{on } W_0^+ - (N(z^f) \times B_+^0); \\ \omega & \text{on } N(z^f) \times B_+^0; \end{cases}$$

and ω is an area form on Σ which agrees with $d_2\beta_t$ on $\Sigma - N(z^f)$. The 2-form Ω_0^+ is symplectic by an easy calculation which uses (f).

Step 4 (Construction of (W_1^+, Ω_1^+) and primitives Θ_0^+, Θ_1^+). Let

$$W_1^+ := S'_{1/2} \times D^2, \quad S'_{1/2} := S_{1/2} - A_{[0,1/2]}.$$

We use polar coordinates (r_2, θ_2) on $D^2 = \{r_2 \leq 1\}$. We identify neighborhoods of $\{0\} \times S'_{1/2} \times \mathbb{R}/2\mathbb{Z} \subset \partial W_0^+$ and $S'_{1/2} \times \partial D^2 \subset \partial W_1^+$ as follows:

$$\phi_{01} : [-\varepsilon', \varepsilon'] \times S'_{1/2} \times \mathbb{R}/2\mathbb{Z} \xrightarrow{\sim} S'_{1/2} \times \{(r_2, \theta_2) \mid e^{-\pi\varepsilon'} \leq r_2 \leq e^{\pi\varepsilon'}\},$$
$$(s, x, t) \mapsto (x, e^{\pi s}, \pi t),$$

where $\varepsilon' > 0$ is sufficiently small.

Let ω_{D^2} be an area form on D^2 satisfying:

$$\omega_{D^2} = \begin{cases} r_2 dr_2 d\theta_2 & \text{near } r_2 = 0; \\ \frac{1}{\pi^2 r_2} dr_2 d\theta_2 & \text{near } r_2 = 1. \end{cases}$$

We then define

$$\Omega_1^+ := \widetilde{\omega}|_{S'_{1/2}} + \omega_{D^2}.$$

An easy calculation shows that $\omega_{D^2} = ds \wedge dt$ (and hence $\Omega_1^+ = \Omega_0^+$) on their overlap.

We write $\omega_{D^2} = d(\phi(r_2) d\theta_2)$, where $\phi : [0,1] \to \mathbb{R}$ satisfies

$$\phi(r_2) = \begin{cases} r_2^2/2 & \text{near } r_2 = 0; \\ \frac{1}{\pi^2} \log r_2 + \frac{1}{10} & \text{near } r_2 = 1. \end{cases}$$

Then $\phi(r_2) d\theta_2 = (s + \frac{\pi}{10}) dt$ on their overlap. The choice of the constant $\frac{\pi}{10} < 1$ will be used in the proof of Lemma 5.4.2. We then define primitives Θ_i^+ of Ω_i^+, $i = 0, 1$, as follows:

$$\Theta_0^+ = (s + \pi/10) dt + \lambda_{+,s} \text{ on } W_0^+ - (N(z^f) \times B_+^0); \tag{4.1.1}$$

$$\Theta_1^+ = \phi(r_2) d\theta_2 + \lambda_{+,0} \text{ on } W_1^+ - (N(z^f) \times D^2). \tag{4.1.2}$$

We have $\Theta_0^+ = \Theta_1^+$ on their overlap.

Step 5 (Corner smoothing). We now have a 4-manifold $W_0^+ \cup W_1^+$ with a concave corner along $(\partial S'_{1/2}) \times \partial D^2$. The component $\mathcal{B}$ of $\partial(W_0^+ \cup W_1^+)$ that contains the corner is homeomorphic to M and $(\partial S'_{1/2}) \times D^2$ is a neighborhood of the binding $(\partial S'_{1/2}) \times \{0\}$.

In this step we round the corner of $\mathcal{B}$ to obtain the smoothing $\widetilde{\mathcal{B}} \subset W_0^+ \cup W_1^+$. We write $\widetilde{\mathcal{B}}_i = \widetilde{\mathcal{B}} \cap W_i^+$, $i = 0, 1$. We define the contact form λ_- on $\widetilde{\mathcal{B}}$ so that

$\lambda_-|_{\widetilde{\mathcal{B}}_i} = \Theta_i^+|_{\widetilde{\mathcal{B}}_i}$, $i = 0, 1$. Here the notation $|_A$ refers to the pullback to A. See Figure 3.


0
1
r_2
r_1
k_0
$\frac{1}{2}+\varepsilon_0$
$\frac{1}{2}$
$\frac{1}{2}-\varepsilon_0$
0
$\widetilde{\mathcal{B}}_1$
W_1^+
W_0^+
W_0^+
$\mathcal{B}$
$\widetilde{\mathcal{B}}_{01}$
$\widetilde{\mathcal{B}}_{00}$
s
0 ε_1

FIGURE 3. Schematic diagram for rounding the corner of $\mathcal{B}$. The diagram shows a neighborhood $N(\mathcal{B})$ of $\mathcal{B}$, where we are projecting $W_0^+ \cap N(\mathcal{B})$ to coordinates (s, r_1) and $W_1^+ \cap N(\mathcal{B})$ to coordinates (r_0, r_1).

Construction of $\widetilde{\mathcal{B}}_0$. There exist $\varepsilon_0, \varepsilon_1 > 0$ small with $\frac{\varepsilon_1}{2\varepsilon_0} < \delta$ and $\varepsilon_1 < \varepsilon'$ and a smooth map $\psi : [0, \frac{1}{2}+\varepsilon_0] \to \mathbb{R}$ such that:

- $\psi(r_1) = \varepsilon_1$ on $[0, \frac{1}{2}-\varepsilon_0]$;
- $-\delta \leq \psi'(r_1) \leq 0$ on $[\frac{1}{2}-\varepsilon_0, \frac{1}{2}+\varepsilon_0]$; and
- $\psi(\frac{1}{2}+\varepsilon_0) = 0$ and $\psi'(\frac{1}{2}+\varepsilon_0) = -\delta$.

We then let $\widetilde{\mathcal{B}}_0 = \widetilde{\mathcal{B}}_{00} \cup \widetilde{\mathcal{B}}_{01}$, where

$$\widetilde{\mathcal{B}}_{00} = \{s = \varepsilon_1\} \times N_{(S_0, \hbar_0)}$$
$$\widetilde{\mathcal{B}}_{01} = \{s = \psi(r_1), r_1 \in [0, 1/2 + \varepsilon_0]\} \times \mathbb{R}/2\mathbb{Z} \times S^1.$$

Here $\mathbb{R}/2\mathbb{Z} \times S^1$ has coordinates (t, θ_1).

Lemma 4.1.3. *There exists* $r_1^* \in (0, \frac{1}{2}+\varepsilon_0)$ *such that each orbit in* $\widetilde{\mathcal{B}}_{01} \cap \{r_1 \neq r_1^*\}$ *is directed by some* $\partial_t + \delta' \partial_{\theta_1}$*, where* $0 < \delta' \leq \delta$ *or* $-\delta \leq \delta' < 0$*, and* $\widetilde{\mathcal{B}}_{01} \cap \{r_1 = 1 + \varepsilon_0\}$ *is directed by* $\partial_t + \delta \partial_{\theta_1}$*.*

Proof. The 1-form $\lambda_-|_{\widetilde{\mathcal{B}}_{00}}$ is clearly a contact form and

$$\lambda_-|_{\widetilde{\mathcal{B}}_{01}} = (\psi(r_1) + f_{0,t}(r_1, \theta_1) + \pi/10)dt + (C + r_1)d\theta_1, \tag{4.1.3}$$

with respect to coordinates (r_1, θ_1, t). The Reeb vector field R_{λ_-} is parallel to $\partial_t - \frac{\partial}{\partial r_1}(\psi + f_{0,t})\partial_{\theta_1}$. Let $r_1^* \in [0, \frac{1}{2}+\varepsilon_0]$ be the point where $\frac{\partial}{\partial r_1}(\psi + f_{0,t}) = 0$.

Then $0 < -\frac{\partial}{\partial r_1}(\psi + f_{0,t}) \leq \delta$ for $r_1 \in [r_1^*, \frac{1}{2} + \varepsilon_0]$, $-\frac{\partial}{\partial r_1}(\psi + f_{0,t})(\frac{1}{2} + \varepsilon_0) = \delta$, and $0 < \frac{\partial}{\partial r_1}(\psi + f_{0,t}) \leq \delta$ for $r_1 \in (0, r_1^*)$, which imply the lemma. $\square$

Construction of $\widetilde{\mathcal{B}}_1$. Let $\zeta : [0,1] \to \mathbb{R}$ be a smooth map such that:

- $\zeta(r_2) = k_0 - k_1 r_2^2/2$ near $r_2 = 0$, where $k_0, k_1 \gg 0$;
- $\zeta'' < 0$ on $(0,1]$;
- $\zeta(1) = \frac{1}{2} + \varepsilon_0$.

We then define $\widetilde{\mathcal{B}}_1 = \{r_1 = \zeta(r_2)\}$.

Lemma 4.1.4. *There exist* $k_0, k_1 \gg 0$, $N = N(k_0, k_1) \gg 0$, *and* ζ *such that* $R_{\lambda_-}|_{\widetilde{\mathcal{B}}_1}$ *is directed by* $\pi\partial_{\theta_2} + \delta\partial_{\theta_1}$.

Proof. $\lambda_-|_{\widetilde{\mathcal{B}}_1}$ is given by

$$\lambda_-|_{\widetilde{\mathcal{B}}_1} = (\phi(r_2) + (1+\varepsilon)/\pi)d\theta_2 + (C + \zeta(r_2))d\theta_1, \tag{4.1.4}$$

with respect to coordinates $(\theta_1, r_2, \theta_2)$. The Reeb vector field R_{λ_-} is parallel to $\pi\partial_{\theta_2} - \frac{\phi'}{\zeta'}\partial_{\theta_1}$. By choosing $k_0, k_1 \gg 0$, $N(k_0, k_1) \gg 0$, and ζ suitably, we may assume that $-\frac{\phi'}{\zeta'}(r_2) = \delta$ for all $r_2 \in (0,1]$. $\square$

We also define $N(K) \subset \widetilde{\mathcal{B}}$ as the closed neighborhood of the binding $K = \{r_2 = 0\}$ that is bounded by the torus $\{r_1 = r_1^*\}$. The region $\mathcal{N} = \{0 < r_1 < r_1^*\} \subset \widetilde{\mathcal{B}}$ will be called "no man's land".

Step 6 (Construction of (W_2^+, Ω_2^+)). Let $W_{01}^+ \subset W_0^+ \cup W_1^+$ be the closure of the component of $(W_0^+ \cup W_1^+) - \widetilde{\mathcal{B}}$ that does not contain $\mathcal{B}$. We then glue the negative cylindrical end

$$(W_2^+, \Omega_2^+) := ((-\infty, 0] \times \widetilde{\mathcal{B}}, d(e^s\lambda_-))$$

to W_{01}^+ along $\widetilde{\mathcal{B}}$, where s is the coordinate for $(-\infty, 0]$. This concludes the construction of (W^+, Ω^+).

4.2. **Further definitions.**

Hamiltonian structure on $\Sigma \times [0,1]$. Let $\overline{\omega} = \widetilde{\omega}|_{s=\frac{3}{2}}$. Then the Hamiltonian structure on $\Sigma \times [0,1]$ at the positive end of W^+ is given by $(dt, \overline{\omega}|_{\Sigma\times[0,1]})$. Let $\hslash_2^+$ be the flow of the corresponding Hamiltonian vector field from $\Sigma \times \{0\}$ to $\Sigma \times \{1\}$. Note that we do not necessarily have $\hslash_2^+ = id$ by construction.

Lagrangian submanifold $L_{\boldsymbol{\alpha}}$. As in Section I.5.2.1, we define the Lagrangian submanifold $L_{\boldsymbol{\alpha}} \subset \partial W^+$ by placing a copy of $\boldsymbol{\alpha}$ on the fiber $\pi^{-1}(3,1)$ over $(3,1) \in \partial B_+^0$ and using the symplectic connection Ω^+ to parallel transport $\boldsymbol{\alpha}$ along the boundary component $(\partial B_+^0) \cap \{s \geq 1\}$ of B_+^0. Observe that

$$L_{\boldsymbol{\alpha}} \cap \{s \geq 3\} = ([3,\infty) \times \{0\} \times \hslash^+ \circ (\hslash_2^+)^{-1}(\boldsymbol{\alpha})) \cup ([3,\infty) \times \{1\} \times \boldsymbol{\alpha}). \tag{4.2.1}$$

Lemma 4.2.1. $\boldsymbol{\beta}' := \hslash^+ \circ (\hslash_2^+)^{-1}(\boldsymbol{\alpha})$ *is isotopic to* $\boldsymbol{\beta}$.

Proof. First observe that $\mathfrak{h}_1^+$ and $\mathfrak{h}_2^+$ are isotopic to the identity. Then $\mathfrak{h}^+$ is isotopic to $\mathfrak{h}_0^+$ where $\mathfrak{h}_0^+|_{S_{1/2}} = id$ and $\mathfrak{h}_0^+|_{S_0}$ is isotopic to $\mathfrak{h}$. The lemma then follows. □

Submanifolds S_z, C_θ, and $\mathcal{H}$. Given $z \in N(z^f)$, let

$$S_z = \{z\} \times (B_+^0 \cup D^2),$$

where $\{z\} \times B_+^0 \subset W_0^+$ and $\{z\} \times D^2 \subset W_1^+$. Also let

$$C_\theta = (\{\theta\} \times B_+^0) \cup (\{\theta\} \times (-\infty, 0] \times \mathbb{R}/2\mathbb{Z}),$$

where $\theta \in \partial S_0$, and let $\mathcal{H} = \cup_{\theta \in \partial S_0} C_\theta$.

Definition of W_+. Let W_+ be the closure of the component of $W^+ - \mathcal{H}$ which is disjoint from $S_{(z')^f}$. In particular, the restriction $\pi_1 : W_+ \cap W_0^+ \to B_+^0$ is a fibration with fiber S_0, $W_+ \cap W_1^+ = \varnothing$, and $W_+ \cap W_2^+ = [0, \infty) \times N_{(S_0, \mathfrak{h}_0)}$. The cobordism W_+ is diffeomorphic to the cobordism used to define the map Φ in Section I.5.1.

4.3. **Proof of Lemma 4.1.1.** (1), (5), (6), (8) are clear from the construction.

(2) follows by letting $\Theta^+ = \Theta_i^+$, $i = 0, 1, 2$, where defined.

(3) By construction, $L_{\boldsymbol{\alpha}}$ is Lagrangian and $d\Theta^+|_{L_{\boldsymbol{\alpha}}} = 0$. It then suffices to observe that $\Theta^+ = 0$ on $L_{\boldsymbol{\alpha}} \cap \pi^{-1}(3, 1)$. This follows from the fact that $\boldsymbol{\alpha} \times \{1\}$ is a Legendrian submanifold of $(N_{(\Sigma - N(z^f), \mathfrak{h}_0^+)}, \lambda_+)$.

(4) The first sentence follows from the construction and the second sentence follows from Lemma 4.2.1.

(7) By Lemma 4.1.4, the Reeb vector field R_{λ_-} has no closed orbits in $\widetilde{\mathcal{B}}_1$ since $\delta > 0$ is irrational. By Lemma 4.1.3 and Equation (4.1.3), each orbit of R_{λ_-} in $\widetilde{\mathcal{B}}_{01} \cap \{r_1 \neq r_1^*\}$ has λ_--action $\geq \frac{1}{2\delta} - (C + \frac{1}{2})$, where $C > 0$ is independent of δ. The second sentence of (7) is immediate from the construction of λ_-.

5. THE CHAIN MAP Φ^+

The goal of this section is to define the chain map

$$\Phi^+ : CF^+(\Sigma, \boldsymbol{\alpha}, \boldsymbol{\beta}, z^f) \to ECC(M, \lambda_-),$$

which is induced by the symplectic cobordism (W^+, Ω^+) and an admissible almost complex structure J^+. We take $\boldsymbol{\beta} = \mathfrak{h}_2^+ \circ \mathfrak{h}^+ \circ (\mathfrak{h}_2^+)^{-1}(\boldsymbol{\alpha})$, in view of Equation (4.2.1) and Lemma 4.2.1 and the fact that $\mathfrak{h}_2^+$ is the flow of the Hamiltonian vector field of $\widetilde{\omega}|_{s=s_0}$, $s_0 \gg 0$, from $\Sigma \times \{0\}$ to $\Sigma \times \{1\}$.

For simplicity we identify $W^+ \cap \{s \geq s_0\} \simeq [s_0, \infty) \times [0, 1] \times \Sigma$ with coordinates (s, t, x) so that $\mathfrak{h}_2^+ = id$ and the Hamiltonian vector field is ∂_t.

5.1. **Almost complex structures.** Let $\overline{\omega} = \widetilde{\omega}|_{s=3/2}$.

Lemma 5.1.1. *There exists a sequence $(\overline{\lambda}_\tau, \overline{\omega})$, $\tau \in [0, 1]$, of stable Hamiltonian structures on $N_{(S_0, \mathfrak{h}_0)}$ such that $\overline{\lambda}_1 = \lambda$, $\overline{\lambda}_\tau$ is a contact form for $\tau > 0$, and $\overline{\lambda}_0 = dt$. The 1-forms $\overline{\lambda}_\tau = f_{t,\tau} dt + \beta_{t,\tau}$ can be normalized so that $\frac{1}{2} < |f_{t,\tau}| \leq 2$.*

Proof. Follows from the discussion of Section I.3.1. □

Definition 5.1.2. An almost complex structure J^+ on W^+ is (W^+,Ω^+)-*admissible* if the following hold:

1. J^+ is tamed by Ω^+;
2. J^+ is s-invariant for $\{s \geq \frac{3}{2}\} \cap W_0^+$ and is adapted to the stable Hamiltonian structure $(dt, \overline{\omega}|_{\Sigma\times[0,1]})$ at the positive end;
3. J^+ is s-invariant for $\{s \leq -\frac{1}{2}\} \cap W_2^+$ and is adapted to the contact form λ_- at the negative end;
4. the restriction J_+ of J^+ to W_+ is C^l-close to a regular admissible almost complex structure J_+^0 on W_+ with respect to $(\overline{\lambda}_0, \overline{\omega})$ (cf. Definitions I.5.4.1 and I.5.8.5);
5. the surfaces $S_{(z')^f}$ and C_θ are J^+-holomorphic for all $\theta \in \partial S_0$.

Let J, J' be the adapted almost complex structures that agree with J^+ at the positive and negative ends.

Note that (4) imposes additional conditions on Ω^+ and λ_-. In practice, the order in which we construct Ω^+ and J^+ is a little convoluted: (i) choose a regular J_+^0, (ii) choose $\tau > 0$ sufficiently small and J_+ sufficiently close to J_+^0, (iii) construct Ω^+ using $\overline{\lambda}_\tau$ in place of λ, and (iv) extend J^+ to the rest of W^+.

Let $\mathcal{J}^+$ be the set of all (W^+,Ω^+)-admissible almost complex structures.

5.2. **The ECH index.**

Let $\mathcal{P} = \mathcal{P}_{\lambda_-}$ be the set of simple orbits of R_{λ_-} and let $\mathcal{O} = \mathcal{O}_{\lambda_-}$ be the set of orbit sets constructed from $\mathcal{P}$.

Let $J^+ \in \mathcal{J}^+$ be an admissible almost complex structure. Let $\mathcal{M}_{J^+}(\mathbf{y},\gamma)$ be the set of holomorphic maps $u : (F,j) \to (W^+, J^+)$ from $\mathbf{y} \in \mathcal{S}_{\boldsymbol{\alpha},\boldsymbol{\beta}}$ to $\gamma \in \mathcal{O}$, such that ∂F is mapped to a distinct component of $L_{\boldsymbol{\alpha}}$ and each component is used exactly once. Elements of $\mathcal{M}_{J^+}(\mathbf{y},\gamma)$ will be called W^+-*curves.*

Let $\check{W}^+$ be W^+ with the ends $\{s > 3\}$ and $\{s < -1\}$ removed and let

$$Z_{\mathbf{y},\gamma} = (L_{\boldsymbol{\alpha}} \cap \check{W}^+) \cup (\{3\} \times [0,1] \times \mathbf{y}) \cup (\{-1\} \times \gamma)$$

as in Section I.5.4.2. The class $[u]$ of $u \in \mathcal{M}_{J^+}(\mathbf{y},\gamma)$ is the relative homology class of the compactification $\check{u}$ in $H_2(\check{W}^+, Z_{\mathbf{y},\gamma})$. Given $A \in H_2(\check{W}^+, Z_{\mathbf{y},\gamma})$, we write $\mathcal{M}_{J^+}(\mathbf{y},\gamma,A) \subset \mathcal{M}_{J^+}(\mathbf{y},\gamma)$ for the subset of W^+-curves u in the class A.

Definition 5.2.1 (Filtration $\mathcal{F}$)**.** Given a W^+-curve u, we define

$$\mathcal{F}(u) = \langle [u], S_{(z')^f} \rangle,$$

where $\langle , \rangle$ is the algebraic intersection number. Since $S_{(z')^f}$ is a holomorphic divisor, $\mathcal{F}(u) \geq 0$.

The definition of the ECH index given in Section I.5.6 also extends directly to our case. The ECH index of a W^+-curve from $\mathbf{y}$ to γ in the class A is denoted by $I_{W^+}(\gamma, A)$.

5.3. **Homology of W^+.**

The goal of this subsection is to compute $H_2(W^+)$. We introduce some notation which will be used only in this subsection: $N = N_{(S_0,\hbar_0)}$, $N_0 = N_{(S_{1/2},\hbar_0^+|_{S_{1/2}})}$ and $\overline{N} = N_{(\Sigma,\hbar_0^+)}$.

Lemma 5.3.1. *$H_2(N) \cong H_2(M)$ and $H_1(N) \cong H_1(M) \oplus \mathbb{Z}$, where the extra $\mathbb{Z}$ factor is generated by a meridian of the binding.*

Proof. The lemma follows from the exact sequence of the pair (M, N). □

Lemma 5.3.2. $H_2(W_0^+) \cong H_2(N) \oplus H_2(N_0) \oplus H_2(\Sigma)$.

Proof. Observe that W_0^+ is homotopy equivalent to $\overline{N}$. We compute $H_2(\overline{N})$ using the Mayer-Vietoris sequence:

$$H_2(N \cap N_0) \xrightarrow{i} H_2(N) \oplus H_2(N_0) \to H_2(\overline{N}) \to H_1(N \cap N_0) \xrightarrow{j} H_1(N) \oplus H_1(N_0).$$

Since $i = 0$ and $\ker j = \mathbb{Z}\langle \partial S_0 \rangle = \mathbb{Z}\langle \partial S_{1/2} \rangle$, the lemma follows. □

Lemma 5.3.3. $H_2(W^+) \cong H_2(M) \oplus H_2(\Sigma)$.

Proof. W^+ is homotopy equivalent to $W_0^+ \cup W_1^+$ and $W_0^+ \cap W_1^+ \cong N_0$. Since W_1^+ is homotopy equivalent to $S_{1/2}$, the Mayer-Vietoris sequence becomes:

$$H_2(N_0) \xrightarrow{i} H_2(W_0^+) \to H_2(W^+) \to H_1(N_0) \xrightarrow{j} H_1(W_0^+) \oplus H_1(S_{1/2}).$$

The map i surjects onto the factor $H_2(N_0)$ in the decomposition of $H_2(W_0^+)$ coming from Lemma 5.3.2. The map j is injective, since $H_1(N_0) \cong H_1(S_{1/2}) \oplus H_1(S^1)$ by the Künneth formula, the restriction $j : H_1(S_{1/2}) \to H_1(S_{1/2})$ is an isomorphism, and the restriction $j : H_1(S^1) \to H_1(\overline{N}) \simeq H_1(W_0^+)$ is injective because the image of the generator of $H_1(S^1)$ is dual to the fiber Σ. The lemma then follows from Lemma 5.3.1. □

5.4. **Energy bound.**

Definition 5.4.1. Let $\mathcal{C}_+$ be the set of nondecreasing functions $[0, +\infty) \to [0, 1]$ and let $\mathcal{C}_-$ be the set of nondecreasing functions $(-\infty, 0] \to [0, 1]$. Let

$$\Omega^+_{\phi,\psi} := \begin{cases} \widetilde{\omega} + d\phi(s) \wedge dt & \text{on } W_0^+ \cap W_{01}^+; \\ \Omega_1^+ & \text{on } W_1^+ \cap W_{01}^+; \\ d(\psi(s)\lambda_-) & \text{on } W_2^+, \end{cases}$$

where $(\phi, \psi) \in \mathcal{C}_+ \times \mathcal{C}_-$.[6] Then the *energy* of a W^+-curve $u : F \to W^+$ from $[\mathbf{y}, i]$ to γ is given by:

$$E(u) = \sup_{\phi,\psi} \int_F u^* \Omega^+_{\phi,\psi}, \tag{5.4.1}$$

where the supremum is taken over all pairs $(\phi, \psi) \in \mathcal{C}_+ \times \mathcal{C}_-$ such that $\Omega^+_{\phi,\psi}$ is smooth.[7]

The condition imposed on the intersection with $S_{(z')^f}$ gives an energy bound:

Lemma 5.4.2 (Energy bound)**.** *For all $k \in \mathbb{N}$, there exists $N_k > 0$ such that $E(u) \leq N_k$ for all $\mathbf{y} \in \mathcal{S}_{\boldsymbol{\alpha},\boldsymbol{\beta}}$, $\gamma \in \mathcal{O}$, and $u \in \mathcal{M}^{\mathcal{F}=k}_{J^+}(\mathbf{y}, \gamma)$.*

---

[6] ϕ, ψ used here are not to be confused with ϕ, ψ which appeared in Section 4.1.

[7] Recall that Ω_1^+ is given by Equation (4.1.2) and agrees with Ω_0^+, given by Equation (4.1.1), on their overlap. This imposes conditions on (ϕ, ψ).

Proof. Let $u : (F, j) \to (W^+, J^+)$ be a holomorphic map in $\mathcal{M}^{\mathcal{F}=k}_{J^+}(\mathbf{y}, \gamma)$. By Lemma 4.1.1(2),(3), $\Omega^+ = d\Theta^+$ on $W^\circ := W^+ - N(S_{zf})$ and the Lagrangian $L_{\boldsymbol{\alpha}}$ is Θ^+-exact. Hence $\int_{\partial F} u^*\Theta^+$ only depends on $\mathbf{y}$.

Let $v : F' \to W^\circ$ be a representative of the homology class $[u] - k[\Sigma] \in H_2(\check{W}^+, Z_{\mathbf{y},\gamma})$. Since the energy is obtained by integrating a closed form,

$$E(u) = E(v) + k \int_\Sigma \widetilde{\omega}.$$

Now $\Omega^+_{\phi,\psi} = d\Theta^+_{\phi,\psi}$ on W°, where

$$\Theta^+_{\phi,\psi} = \begin{cases} \lambda_{+,s} + \phi(s)dt & \text{on } W_0^+ \cap W^\circ \cap W_{01}^+; \\ \Theta_1^+ & \text{on } W_1^+ \cap W^\circ \cap W_{01}^+; \\ \psi(s)\lambda_- & \text{on } W_2^+. \end{cases}$$

By Equations (4.1.1) and (4.1.2), Θ_1^+ can be written as $\lambda_{+,s} + (s + \frac{\pi}{10})dt$ on $W_0^+ \cap W_1^+ \cap W^\circ \cap W_{01}^+$. Since $\frac{\pi}{10} < 1$, there exist smooth $\Theta^+_{\phi,\psi}$ that extend Θ_1^+, i.e., there exist $\phi \in \mathcal{C}_+$ that extend $\phi(s) = s + \frac{\pi}{10}$ near $s = 0$.

By Stokes' theorem,

$$\begin{aligned} E(v) &\leq \int_{\{s\}\times[0,1]\times\mathbf{y}} \lambda_+ + \sup_{\phi\in\mathcal{C}_+} \lim_{s\to\infty} \int_{\{s\}\times[0,1]\times\mathbf{y}} \phi dt \\ &\quad + \int_{\partial F} v^*\lambda_+ + \sup_{\phi\in\mathcal{C}_+} \int_{\partial F} \phi dt - \inf_{\psi\in\mathcal{C}_-} \int_\gamma \psi\lambda_- \\ &\leq 4g + \int_{[0,1]\times\mathbf{y}} \lambda_+ + \int_{\partial F} v^*\lambda_+. \end{aligned} \tag{5.4.2}$$

Recall that $\lambda_{+,s} = \lambda_+$ for $s \geq \frac{3}{2}$. In the above calculation,

$$\sup_{\phi\in\mathcal{C}_+} \lim_{s\to\infty} \int_{\{s\}\times[0,1]\times\mathbf{y}} \phi dt = 2g, \ \sup_{\phi\in\mathcal{C}_+} \int_{\partial F} \phi dt = 2g, \ \inf_{\psi\in\mathcal{C}_-} \int_\gamma \psi\lambda_- = 0.$$

We then obtain

$$E(u) \leq 4g + \int_{[0,1]\times\mathbf{y}} \lambda_+ + \int_{\partial F} u^*\lambda_+ + k\int_\Sigma \widetilde{\omega},$$

which is the desired bound. □

5.5. **Regularity.** Define the subset $\mathcal{M}^h_{J^+}(\mathbf{y}, \gamma, A) \subset \mathcal{M}_{J^+}(\mathbf{y}, \gamma, A)$ consisting of holomorphic curves without vertical fiber components. As in Lemma I.5.8.2, the set $\mathcal{J}^+_{reg}$ of *regular* $J^+ \in \mathcal{J}^+$ for which all the moduli spaces $\mathcal{M}^h_{J^+}(\mathbf{y}, \gamma, A)$ are transversally cut out is a dense subset of $\mathcal{J}^+$. We can restrict attention to $\mathcal{M}^h_{J^+}(\mathbf{y}, \gamma, A)$ for the following reason:

Lemma 5.5.1. *If $J^+ \in \mathcal{J}^+_{reg}$ and $u \in \mathcal{M}_{J^+}(\mathbf{y}, \gamma, A) - \mathcal{M}^h_{J^+}(\mathbf{y}, \gamma, A)$, then $I_{W^+}(u) \geq 2 + 2g$.*

Proof. Suppose $u = u_1 \cup u_2$, where u_1 is regular and u_2 is homologous to $k \geq 1$ times a fiber. Since $\langle u_1, u_2 \rangle = k \cdot 2g$,

$$\begin{aligned} I(u) &= I(u_1) + I(u_2) + 2k(2g) \\ &\geq 0 + k(2-2g) + 4kg \geq k(2+2g). \end{aligned}$$

Here $I(u_1) \geq 0$ since $I(u_1) \geq \operatorname{ind}(u_1)$ by the index inequality and $\operatorname{ind}(u_1) \geq 0$ by the regularity of u_1. □

5.6. **Holomorphic curves in W^+ without positive ends.**

In this subsection and the next, we make essential use of the assumption $g(S) \geq 2$.

Let $S'' = S_{1/2} - A_{[0,N]}$ and $\overline{S}'' = S'' \cup \{\infty\}$. We define the "projection" $\pi_{\overline{S}''} : W^+ \to \overline{S}''$ as follows:

- on W_0^+, $\pi_{\overline{S}''}(s,x,t) = x$ if $x \in S''$ and $\pi_{\overline{S}''}(s,x,t) = \infty$ if $x \notin S''$;
- on W_1^+, $\pi_{\overline{S}''}(x,r_2,\theta_2) = x$ if $x \in S''$ and $\pi_{\overline{S}''}(x,r_2,\theta_2) = \infty$ if $x \notin S''$;
- $\pi_{\overline{S}''}(W_2^+) = \infty$.

Lemma 5.6.1. *If $u : \dot{F} \to (W^+, J^+)$ is a holomorphic map without positive ends, then $g(F) \geq 2$.*

Proof. The map $\pi_{\overline{S}''} \circ u$ can be extended to a continuous map $f : F \to \overline{S}''$. Observe that the curve u must intersect $S_{(z')^f}$ because the symplectic form is exact on $W^+ - S_{(z')^f}$. Hence $\deg f > 0$. Now we use the following fact: If $f : \Sigma_1 \to \Sigma_2$ is a positive degree map between closed oriented surfaces, then $g(\Sigma_1) \geq g(\Sigma_2)$. Since $g(S) = g(\overline{S}'') \geq 2$, it follows that $g(F) \geq 2$. □

Lemma 5.6.2. *There are no $I = 0$ closed holomorphic curves in (W^+, J^+).*

Proof. We argue by contradiction. Let $A = [u_*(F)]$. By Lemma 5.3.3, the intersection form on $H_2(W^+)$ is trivial. Hence $A \cdot A = 0$. If $I(A) = A \cdot A + c_1(A) = 0$, then it follows that $c_1(A) = 0$.

Suppose that u is simple. Then $\chi(F) \geq 0$ by the adjunction formula. This contradicts Lemma 5.6.1. In particular $I(u) > 0$ by the regularity of u and the index inequality. If v is a degree d branched cover of u in the class A, then $I(v) = I(dA) = dI(A) \geq d$ using the formula

$$I(dA) = dI(A) + (d^2 - d)A \cdot A. \tag{5.6.1}$$

□

Lemma 5.6.3. *A multiply-covered holomorphic curve u with only negative ends has $I(u) > 0$.*

Proof. This follows from Lemma 5.3.3 and the inequality

$$I(dC) \geq dI(C) + \frac{(d^2-d)}{2}(2g(C) - 2 - \operatorname{ind}(C) + h) \tag{5.6.2}$$

from [Hu2, Section 5.1], where C is a simple curve, $\operatorname{ind}(C)$ is the Fredholm index of C, and h is the number of hyperbolic ends. □

5.7. **The map Φ^+.** Let $J^+ \in \mathcal{J}^+_{reg}$. The chain map Φ^+ is given as follows:

$$\Phi^+ : (CF^+(\Sigma, \boldsymbol{\alpha}, \boldsymbol{\beta}, z^f), \partial) \to (ECC(M, \lambda_-), \partial'),$$
$$[\mathbf{y}, i] \mapsto \sum_{\gamma, A} \#\mathcal{M}^{\mathcal{F}=i, I_{W^+}=0}_{J^+}(\mathbf{y}, \gamma, A) \cdot \gamma,$$

where the summation is over all $\gamma \in \mathcal{O}_{\lambda_-}$ and $A \in H_2(\check{W}^+, Z_{\mathbf{y},\gamma})$. Here ∂' is the usual ECH differential on $ECC(M, \lambda_-)$.

By a combination of Lemma 5.4.2 and the Gromov-Taubes compactness theorem (cf. Section I.3.4), the sum in the definition of Φ^+ is finite. Hence Φ^+ is well-defined.

Theorem 5.7.1. *If $g(S) \geq 2$, then Φ^+ is a chain map.*

Proof. Similar to that of Theorem I.6.2.4, with slight modifications in view of Lemmas 5.6.2 and 5.6.3. □

Remark 5.7.2. One can define the twisted coefficient analog of Φ^+, taking into account Lemma 5.3.3.

5.8. **Restriction to Φ.** In this subsection δ still denotes the constant that appears in the construction of λ_-. Let $\mathcal{P}|_N$ be the subset of $\mathcal{P}$ consisting of orbits that are contained in N. Also let $\gamma_\theta \in \mathcal{P}_-$ be the orbit corresponding to $\theta \in \partial S_0$.

Lemma 5.8.1. *For $\delta > 0$ sufficiently small, if $u \in \mathcal{M}^{\mathcal{F}=0}_{J^+}(\mathbf{y}, \gamma)$, $\mathbf{y} \in \mathcal{S}_{\boldsymbol{\alpha},\boldsymbol{\beta}}$, $\mathbf{y} \subset S_0$, and $\gamma \in \mathcal{O}$, then γ is constructed from $\mathcal{P}|_N \cup \{e', h'\}$.*

Proof. This is similar to Lemma 5.4.2. Let $u : (F, j) \to (W^+, J^+)$ be a holomorphic map in $\mathcal{M}^{\mathcal{F}=0}_{J^+}(\mathbf{y}, \gamma)$. Since $\mathrm{Im}(u) \cap S_{(z')^f} = \varnothing$, we may assume that $\mathrm{Im}(u) \cap N(S_{z^f}) = \varnothing$ by possibly modifying u in its homology class. By considerations similar to those of Lemma 5.4.2, we obtain:

$$4g + \int_{[0,1]\times\mathbf{y}} \lambda_+ + \int_{\partial F} u^*\lambda_+ \geq \mathcal{A}_{\lambda_-}(\gamma),$$

where $\mathcal{A}_{\lambda_-}(\gamma)$ is the action of γ with respect to λ_-. Since the Lagrangian $L_{\boldsymbol{\alpha}}$ is exact by Lemma 4.1.1(3), we may take $u(\partial F) \subset \partial W_+$. Hence there exist upper bounds for $\int_{\partial F} u^*\lambda_+$ and $\int_{[0,1]\times\mathbf{y}} \lambda_+$ which are independent of $\mathbf{y}$ and δ. By Lemma 4.1.1(7), all the orbit sets γ in $int(N(K)) \cup \mathcal{N}$ satisfy $\mathcal{A}_{\lambda_-}(\gamma) \geq \frac{1}{2\delta} - \kappa$. Hence, for $\delta > 0$ sufficiently small, no negative end of u is asymptotic to an orbit in $int(N(K)) \cup \mathcal{N}$. □

Lemma 5.8.2. *If $u \in \mathcal{M}^{\mathcal{F}=0}_{J^+}(\mathbf{y}, \gamma)$, where $\mathbf{y} \in \mathcal{S}_{\boldsymbol{\alpha},\boldsymbol{\beta}}$, $\mathbf{y} \subset S_0$, and γ is constructed from $\mathcal{P}|_N \cup \{e', h'\}$, then $\mathrm{Im}(u) \subset W_+$ and $\gamma \in \mathcal{O}|_N$.*

Proof. Let $u \in \mathcal{M}^{\mathcal{F}=0}_{J^+}(\mathbf{y}, \gamma)$ such that $u(F) \not\subset W_+$.

Suppose that u is not a multi-level Morse-Bott building. Then $u(F) \cap C_{\theta_0} \neq \varnothing$ for some $\theta_0 \in \partial S_0 - \boldsymbol{\alpha} - \boldsymbol{\beta}$, and moreover we may assume that γ_{θ_0} is not an asymptotic limit of u at $-\infty$. Since J^+ is admissible, all the curves C_θ are holomorphic. Hence $\langle u(F), C_{\theta_0} \rangle > 0$ by the positivity of intersections.

Let D_θ, $\theta\in\partial S_0$, be a meridian disk of the solid torus $\mathcal{N}\cup N(K)$ that is bounded by $\{\theta\}\times\mathbb{R}/2\mathbb{Z}$ and is disjoint from $\{e',h'\}$, and let $D_{\theta,s}=\{s\}\times D_\theta\subset W_2^+$, where $s<0$ and $\theta\in\partial S_0$. We then define

$$C_{\theta,s_0}:=(C_\theta-\{s<s_0\})\cup D_{\theta,s_0},$$

where $s_0<0$. When s_0 is sufficiently negative, the curve $u(F)$ intersects C_{θ_0,s_0} only in the region $C_{\theta_0}-\{s<s_0\}$, since γ is constructed from $\mathcal{P}|_N\cup\{e',h'\}$ and D_{θ_0} does not intersect e' and h'. Hence $\langle u(F),C_{\theta_0,s_0}\rangle>0$. Now, since $[S_{(z')^f}]=[C_{\theta_0,s_0}]$ in $H_2(\check{W}^+,\partial\check{W}^+-Z_{\mathbf{y},\gamma})$, we have

$$\mathcal{F}(u)=\langle[u],S_{(z')^f}\rangle=\langle[u],C_{\theta_0,s_0}\rangle>0.$$

This contradicts our assumption that $\mathcal{F}(u)=0$.

If u is a multi-level Morse-Bott building, then we need to make the appropriate modifications (left to the reader), but the same argument goes through. For example, we need to replace C_{θ_0} by a multi-level building $C_{\theta_0}\cup(\mathbb{R}\times\gamma_{\theta_0})\cup\cdots\cup(\mathbb{R}\times\gamma_{\theta_0})$. Note that if u is a Morse-Bott building, then it could have a component u_1 with a negative end that limits to some γ_{θ_1}, followed by a gradient trajectory from θ_1 to θ_2, and then by a component u_2 with a positive end that limits to γ_{θ_2}. □

Theorem 5.8.3. *For* $\delta>0$ *sufficiently small, if* $u\in\mathcal{M}^{\mathcal{F}=0}_{J^+}(\mathbf{y},\gamma)$, $\mathbf{y}\in\mathcal{S}_{\boldsymbol{\alpha},\boldsymbol{\beta}}$, $\mathbf{y}\subset S_0$, *and* $\gamma\in\mathcal{O}$, *then* $\operatorname{Im}(u)\subset W_+$ *and* $\gamma\in\mathcal{O}|_N$.

Proof. Follows from Lemmas 5.8.1 and 5.8.2. □

The restriction Φ of Φ^+ to (W_+,J_+) is given as follows:

$$\Phi:\widehat{CF}(\Sigma,\boldsymbol{\alpha},\boldsymbol{\beta},z^f)\to ECC_{2g}(M,\lambda_-),$$

$$[\mathbf{y},0]\mapsto\sum_{\gamma,A}\#\mathcal{M}^{I_{W_+}=0}_{J_+}(\mathbf{y},\gamma,A)\cdot\gamma,$$

where $\mathcal{M}^{I_{W_+}=0}_{J_+}(\mathbf{y},\gamma,A)$ is the subset of $\mathcal{M}_{J^+}(\mathbf{y},\gamma,A)$ consisting of curves with image in W_+.

Theorem 5.8.4. Φ *is a quasi-isomorphism.*

Proof. The almost complex structure J_+ is sufficiently close to J_+^0. For J_+^0, the analogous chain map was shown to be a quasi-isomorphism (Theorem II.1.0.1). Considerations similar to those of Theorem I.3.6.1 imply that Φ is a quasi-isomorphism. □

5.9. **Commutativity with the U-map.** Let z^b be a point in $\mathbb{R}\times[0,1]$ with t-coordinate $\frac{1}{2}$ and let $z=(z^b,z^f)\in W$. Let U_z be the geometric U-map with respect to z on the HF side. On the ECH side, let $z'=(s,z^M)$ be a generic point in $\mathbb{R}\times int(N(K))$ near the binding K. We define $U'=U'_{z'}$ so that $\langle U'(\gamma),\gamma'\rangle$ is the count of $I_{ECH}=2$ curves in the symplectization $(\mathbb{R}\times M,J')$ from γ to γ' that pass through z'.

Theorem 5.9.1. *There exists a chain homotopy*

$$K : CF^+(\Sigma, \boldsymbol{\alpha}, \boldsymbol{\beta}, z^f) \to ECC(M, \lambda_-)$$

which satisfies

$$U' \circ \Phi^+ - \Phi^+ \circ U_z = \partial' \circ K + K \circ \partial.$$

Proof. The commutativity of Φ^+ with the U-maps up to homotopy is obtained by moving the point constraint in the cobordism W^+ from $s = +\infty$ to $s = -\infty$.

The 1-parameter family of points $(z_\tau)_{\tau \in \mathbb{R}}$ is chosen as follows: For $\tau \geq 0$, let $z_\tau = (z_\tau^b, z^f)$, where z_τ^b approaches $(s,t) = (+\infty, \frac{1}{2})$ as $\tau \to +\infty$ and z_0^b is near the center of the disk $D^2 = \{r_2 \leq 1\}$. Next, for $\tau \in [-1, 0]$, let $z_\tau = (z_0^b, z_\tau^f)$ so that $(z_0^b, z_{-1}^f) \in \{0\} \times \widetilde{\mathcal{B}}$ is near the binding K. For $\tau \leq -1$, let $z_\tau = (\tau + 1, z^M) \in (-\infty, 0] \times M$, where $z^M \in M = \widetilde{\mathcal{B}}$ is a point near the binding. Finally, we consider a small perturbation of $(z_\tau)_{\tau \in \mathbb{R}}$ to make it generic (without changing its name).

We define the 1-parameter family of almost complex structures $(J_\tau^+)_{\tau \in \mathbb{R}}$ so that J_τ^+ is C^l-close to J^+ and agrees with J^+ outside a small neighborhood of z_τ.

The rest of the chain homotopy argument is standard, with the exception of the obstruction theory that was carried out in [HT1, HT2]. □

Theorem 5.9.2. *For $\delta > 0$ sufficiently small, if $\mathbf{y} \in \mathcal{S}_{\boldsymbol{\alpha}, \boldsymbol{\beta}}$ and $\mathbf{y} \subset S_0$, then $K([\mathbf{y}, 0]) = 0$.*

Proof. Similar to the proof of Theorem 5.8.3. The coefficient $\langle K([\mathbf{y}, 0]), \gamma \rangle$ is given by the count of $I_{W^+} = 1$ curves from $\mathbf{y}$ to γ that pass through z_τ for some τ and do not intersect $S_{(z')^f}$. If such a curve u exists, then $\mathrm{Im}(u) \not\subset W_+$. This is not possible by Theorem 5.8.3. □

6. PROOF OF THEOREM 1.0.1

In this section we prove Theorem 1.0.1. In Section 6.1 we prove an algebraic result (Theorem 6.1.5) which is sufficient to prove that Φ^+ is a quasi-isomorphism. The conditions of Theorem 6.1.5 are verified in Section 6.4.

6.1. **Some algebra.**

Definition 6.1.1. Let (A, d) be a chain complex. We say that a chain map $f \colon A \to A$ is *homologically almost nilpotent* (abbreviated *han*) if for every $x \in H(A)$ there exists $n \in \mathbb{N}$ such that $(f_*)^n(x) = 0$.

Prototypical examples of *han* maps are the U-maps in HF^+ and ECH.

Let (A, d_A) and (B, d_B) be chain complexes with *han* maps $U_A \colon A \to A$ and $U_B \colon B \to B$ and let $\Phi^+ \colon A \to B$ be a chain map such that the diagram

$$\begin{tikzcd} A \arrow[r, "\Phi^+"] \arrow[d, "U_A"'] & B \arrow[d, "U_B"] \\ A \arrow[r, "\Phi^+"] & B \end{tikzcd}$$

commutes up to a chain homotopy K. We form a chain complex $D = A \oplus A \oplus B \oplus B$ with differential

$$d_D = \begin{pmatrix} d_A & 0 & 0 & 0 \\ U_A & d_A & 0 & 0 \\ \Phi^+ & 0 & d_B & 0 \\ K & \Phi^+ & U_B & d_B \end{pmatrix}.$$

Given a chain map f, we denote its mapping cone by $C(f)$.

Lemma 6.1.2. *There is an exact triangle:*

$$\begin{tikzcd} H(C(U_A)) \arrow[rr, "(\Phi_{alg})_*"] & & H(C(U_B)) \arrow[dl] \\ & H(D) \arrow[ul] & \end{tikzcd} \tag{6.1.1}$$

where $\Phi_{alg} = \begin{pmatrix} \Phi^+ & 0 \\ K & \Phi^+ \end{pmatrix}$.

Proof. From the shape of d_D, it is evident that (D, d_D) is the mapping cone of $\Phi_{alg}\colon C(U_A) \to C(U_B)$. □

Lemma 6.1.3. *There is an exact triangle:*

$$\begin{tikzcd} H(C(\Phi^+)) \arrow[rr, "(U_{\Phi^+})_*"] & & H(C(\Phi^+)) \arrow[dl] \\ & H(D) \arrow[ul] & \end{tikzcd} \tag{6.1.2}$$

where $U_{\Phi^+} = \begin{pmatrix} U_A & 0 \\ K & U_B \end{pmatrix}$.

Proof. Let $C(\Phi^+) = A \oplus B$ be the cone of Φ^+ with differential $d_{U(\Phi^+)} = \begin{pmatrix} d_A & 0 \\ \Phi^+ & d_B \end{pmatrix}$. Then $U_{\Phi^+} : (C(\Phi^+), d_{U(\Phi^+)}) \to (C(\Phi^+), d_{U(\Phi^+)})$ is a chain map. Hence the complex $(D', d_{D'})$, where $D' = A \oplus B \oplus A \oplus B$ and

$$d_{D'} = \begin{pmatrix} d_{U(\Phi^+)} & 0 \\ U_{\Phi^+} & d_{U(\Phi^+)} \end{pmatrix} = \begin{pmatrix} d_A & 0 & 0 & 0 \\ \Phi^+ & d_B & 0 & 0 \\ U_A & 0 & d_A & 0 \\ K & U_B & \Phi^+ & d_B \end{pmatrix},$$

is the cone of U_{Φ^+}. Moreover $f\colon D \to D'$ where

$$f = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

is an isomorphism of complexes. □

Lemma 6.1.4. *U_{Φ^+} is a* han *map.*

Proof. Consider the following commutative diagram with exact rows:

$$\begin{tikzcd}
H(B) \arrow[r, "i_*"] \arrow[d, "U_B^n"'] & H(C(\Phi^+)) \arrow[r, "j_*"] \arrow[d, "U_{\Phi^+}^n"'] & H(A) \arrow[d, "U_A^n"'] \\
H(B) \arrow[r, "i_*"] \arrow[d, "U_B^m"'] & H(C(\Phi^+)) \arrow[r, "j_*"] \arrow[d, "U_{\Phi^+}^m"'] & H(A) \arrow[d, "U_A^m"'] \\
H(B) \arrow[r, "i_*"] & H(C(\Phi^+)) \arrow[r, "j_*"] & H(A)
\end{tikzcd}$$

Given $x \in H(C(\Phi^+))$, we choose $n \in \mathbb{N}$ sufficiently large so that $U_A^n(j_*(x)) = j_*(U_{\Phi^+}^n(x)) = 0$. Then $U_{\Phi^+}^n(x) = i_*(y)$ for some $y \in H(B)$. Next choose $m \in \mathbb{N}$ sufficiently large so that $U_B^m(y) = 0$. Then $U_{\Phi^+}^{n+m}(x) = U_{\Phi^+}^m(i_*(y)) = i_*(U_B^m(y)) = 0$. □

Theorem 6.1.5. *If Φ_{alg} is a quasi-isomorphism, then Φ^+ is a quasi-isomorphism.*

Proof. If Φ_{alg} is a quasi-isomorphism, then $H(D) = 0$ by Exact Triangle (6.1.1). This in turn implies that U_{Φ^+} is a quasi-isomorphism by Exact Triangle (6.1.2). However the *han* map U_{Φ^+} cannot be a quasi-isomorphism, unless $H(C(\Phi^+)) = 0$. Finally, the triangle

$$\begin{tikzcd}
H(A) \arrow[rr, "\Phi^+_*"] & & H(B) \arrow[dl] \\
& H(C(\Phi^+)) \arrow[ul] &
\end{tikzcd}$$

implies that Φ^+ is a quasi-isomorphism. □

We finish this subsection with a lemma which compares the homology of $C(U)$ with that of $\ker U$.

Lemma 6.1.6. *Let (C, d) be a chain complex and let U be a chain map. If U is surjective, then the inclusion*

$$i : \ker U \to C(U)$$
$$x \mapsto \begin{pmatrix} x \\ 0 \end{pmatrix}$$

is a quasi-isomorphism.

Proof. Let $\overline{U} : C/\ker U \to C$ be the map induced by U. We have a short exact sequence of complexes

$$0 \to \ker U \to C(U) \to C(\overline{U}) \to 0,$$

which induces the exact triangle:

$$\begin{tikzcd}
H(\ker U) \arrow[rr, "i_*"] & & H(C(U)) \arrow[dl] \\
& H(C(\overline{U})) \arrow[ul] &
\end{tikzcd}$$

Since U is surjective, $\overline{U}$ is an isomorphism. Hence $H(C(\overline{U}))=0$ and the lemma follows. $\square$

6.2. **Heegaard Floer chain complexes.** Recall the subcomplex $\widehat{CF'}(S_0,\mathbf{a},\hbar(\mathbf{a}))$ of $\widehat{CF}(\Sigma,\boldsymbol{\alpha},\boldsymbol{\beta},z^f)$ from Section I.4.9.3, which is generated by $\mathcal{S}_{\mathbf{a},\hbar(\mathbf{a})}$; let

$$j':\widehat{CF'}(S_0,\mathbf{a},\hbar(\mathbf{a}))\to\widehat{CF}(\Sigma,\boldsymbol{\alpha},\boldsymbol{\beta},z^f)$$

be the natural inclusion map. We are viewing

$$\widehat{CF}(\Sigma,\boldsymbol{\alpha},\boldsymbol{\beta},z^f)\subset CF^+(\Sigma,\boldsymbol{\alpha},\boldsymbol{\beta},z^f)$$

as the subcomplex generated by elements of the form $[\mathbf{y},0]$. The chain complex $\widehat{CF}(S_0,\mathbf{a},\hbar(\mathbf{a}))$ is the quotient $\widehat{CF'}(S_0,\mathbf{a},\hbar(\mathbf{a}))/\sim$, defined in Section I.4.9.3.

Lemma 6.2.1. *There is an isomorphism* $j:\widehat{HF}(S_0,\mathbf{a},\hbar(\mathbf{a}))\to\widehat{HF}(\Sigma,\boldsymbol{\alpha},\boldsymbol{\beta},z^f)$ *given by* $[Z]\mapsto[Z]$.

Proof. This follows from the discussion of Theorem I.4.9.4. Note that the natural candidate

$$\widehat{CF}(S_0,\mathbf{a},\hbar(\mathbf{a}))\to\widehat{CF}(\Sigma,\boldsymbol{\alpha},\boldsymbol{\beta},z^f),\quad [Z]\to Z$$

for a chain map is not a well-defined map. $\square$

Lemma 6.2.2. *The inclusion* $i:\widehat{CF}(\Sigma,\boldsymbol{\alpha},\boldsymbol{\beta},z^f)\to C(U)$ *given by* $\mathbf{y}\mapsto\begin{pmatrix}[\mathbf{y},0]\\0\end{pmatrix}$ *is a quasi-isomorphism.*

Proof. This follows from Lemma 6.1.6, since $U([\mathbf{y},i])=[\mathbf{y},i-1]$ for $i\geq 1$ and $\ker U\simeq\widehat{CF}(\Sigma,\boldsymbol{\alpha},\boldsymbol{\beta},z^f)$. $\square$

6.3. **ECH chain complexes.** We describe several ECH chain complexes that are related to $(ECC(M,\lambda_-),\partial')$ and are constructed from certain subsets $\mathcal{S}$ of the set $\mathcal{P}=\mathcal{P}_{\lambda_-}$ of simple orbits of R_{λ_-}. Many of these appeared in [CGH1, Section 10] (some with different names). Let U' be the U-map of $ECC(M,\lambda_-)$ with respect to $(s_0,z^M)\in\mathbb{R}\times M$, where z^M is a generic point which is sufficiently close to the binding.

Let $\mathcal{O}_{\mathcal{S}}$ be the set of orbit sets that are constructed from $\mathcal{S}$. Then $\mathcal{S}$ is *closed* if $\gamma'\in\mathcal{O}_{\mathcal{S}}$, whenever $\gamma\in\mathcal{O}_{\mathcal{S}}$, $\gamma'\in\mathcal{O}_{\mathcal{P}}$, and $\langle\partial'\gamma,\gamma'\rangle\neq 0$ or $\langle U'\gamma,\gamma'\rangle\neq 0$. If $\mathcal{S}$ is closed, then let $(A_{\mathcal{S}},\partial'_{\mathcal{S}})$ be the subcomplex of $ECC(M,\lambda_-)$ generated by $\mathcal{O}_{\mathcal{S}}$ and let $U'_{\mathcal{S}}$ be the restriction of U' to $A_{\mathcal{S}}$. Let $\mathcal{P}|_N\subset\mathcal{P}$ be the set of orbits in the suspension N. The subsets

$\mathcal{S}_1=\mathcal{P}|_N\cup\{e',h'\},\ \mathcal{S}_2=\mathcal{P}|_{N\cup\mathcal{N}}\cup\{e',h'\},\ \mathcal{P}|_N\cup\{h'\},\ \mathcal{P}|_{N\cup\mathcal{N}}\cup\{h'\},\ \mathcal{P}|_N$

are closed and we write $A_i = A_{\mathcal{S}_i}$, $\partial_i' = \partial_{\mathcal{S}_i}'$, and $U_i' = U_{\mathcal{S}_i}'$ for $i = 1, 2$, as well as

$$ECC^{\natural}(N) = A_{\mathcal{P}|_N \cup \{h'\}},\ ECC^{\natural\natural}(N) = A_{\mathcal{P}|_{N\cup\mathcal{N}} \cup \{h'\}},\ ECC(N) = A_{\mathcal{P}|_N}.$$

Also let $ECC_{2g}(N) \subset ECC(N)$ be the subcomplex generated by orbit sets γ satisfying $\langle \gamma, S \times \{t\}\rangle = 2g$. Let

$$q_1 : ECC_{2g}(N) \to ECC^{\natural}(N), \quad q_2 : ECC_{2g}(N) \to ECC^{\natural\natural}(N)$$

be the chain maps given by the natural inclusion. Then we have the following:

Lemma 6.3.1. *The chain maps q_1 and q_2 are quasi-isomorphisms.*

Proof. The chain map q_1 is a quasi-isomorphism by Section II.5 and [CGH1, Section 10.5]. By a direct limit argument similar to that of [CGH1, Proposition 7.2.2], there is a quasi-isomorphism $r : ECC^{\natural\natural}(N) \to ECC^{\natural}(N)$ such that $r \circ q_2 = q_1$. This implies that q_2 is also a quasi-isomorphism. ☐

Lemma 6.3.2. *The inclusions $p_1 : ECC^{\natural}(N) \to C(U_1')$ and $p_2 : ECC^{\natural\natural}(N) \to C(U_2')$ given by $\Gamma \mapsto \begin{pmatrix}\Gamma\\0\end{pmatrix}$ are quasi-isomorphisms.*

Proof. This follows from Lemma 6.1.6. The map U_i', $i = 1, 2$, is given by:

$$U_i'((e')^k (h')^l \Gamma) = (e')^{k-1}(h')^l \Gamma, \tag{6.3.1}$$

where $\Gamma \in \mathcal{O}|_N$ or $\mathcal{O}|_{N\cup\mathcal{N}}$; see [CGH1, Claim 10.5.1] for a similar calculation. Hence U_i' is surjective, $\ker U_1' = ECC^{\natural}(N)$, and $\ker U_2' = ECC^{\natural\natural}(N)$. This implies the lemma. ☐

Lemma 6.3.3. *The inclusion $\mathfrak{i} : ECC^{\natural\natural}(N) \to C(U')$ given by $\Gamma \mapsto \begin{pmatrix}\Gamma\\0\end{pmatrix}$ is a quasi-isomorphism.*

Proof. This is similar to the argument in [CGH1, Section 10].

Choose an identification $\eta : H_1(N(K);\mathbb{Z}) \xrightarrow{\sim} \mathbb{Z}$ such that the homology class of the binding is 1. Define the filtration $\mathcal{F} : ECC(M) \to \mathbb{Z}^{\geq 0}$ such that

$$\mathcal{F}\left(\sum_i \gamma_i \otimes \Gamma_i\right) = \max_i \eta([\gamma_i]),$$

where $\gamma_i \in \mathcal{O}|_{N(K)}$ and $\Gamma_i \in \mathcal{O}|_{N\cup\mathcal{N}}$. Let $\mathcal{F}^{\natural\natural} : ECC^{\natural\natural}(N) \to \mathbb{Z}^{\geq 0}$ be its restriction to $ECC^{\natural\natural}(N)$. (Note that $\mathcal{F}^{\natural\natural}$ is a trivial filtration.) Next define the filtration $\widehat{\mathcal{F}} : C(U') \to \mathbb{Z}^{\geq 0}$ such that

$$\widehat{\mathcal{F}}\begin{pmatrix} \sum_i \gamma_i \otimes \Gamma_i \\ \sum_j \gamma_j' \otimes \Gamma_j' \end{pmatrix} = \max_{i,j}\{\eta([\gamma_i]), \eta([\gamma_j'])\}.$$

The map $\mathfrak{i}$ is an $(\mathcal{F}^{\natural\natural}, \widehat{\mathcal{F}})$-filtered chain map. The induced map

$$E^1(\mathfrak{i}) : E^1(\mathcal{F}^{\natural\natural}) \to E^1(\widehat{\mathcal{F}})$$

on the E^1-level agrees with the isomorphism $(p_2)_*$; the proof is similar to that of [CGH1, Section 10]. If a filtered chain map between filtered chain complexes which are bounded below is an isomorphism on the E^r-level, then it is a quasi-isomorphism. This implies that $\mathfrak{i}$ is a quasi-isomorphism. $\square$

6.4. **Completion of proof of Theorem 1.0.1.** By Theorems 3.1.4, 5.7.1, and 5.9.1, the map

$$\Phi^+ : CF^+(\Sigma, \boldsymbol{\alpha}, \boldsymbol{\beta}, z^f) \to ECC(M, \lambda_-)$$

is a chain map which commutes with U and U' up to the chain homotopy $K^+ = K + \Phi^+ \circ H$, where H is given in Theorem 3.1.4 and K is given in Theorem 5.9.1. Here U is the formal U-map on $(CF^+(\Sigma, \boldsymbol{\alpha}, \boldsymbol{\beta}, z^f), \partial)$ and U' is the U-map on $(ECC(M, \lambda_-), \partial')$. In view of Theorem 6.1.5, Theorem 1.0.1 immediately follows from:

Theorem 6.4.1. *The algebraic map Φ_{alg} is a quasi-isomorphism.*

Let $\Phi' : \widehat{CF}'(S_0, \mathbf{a}, \hbar(\mathbf{a})) \to ECC_{2g}(N)$ be the map from Definition I.6.2.1. The map Φ' descends to $\Phi : \widehat{CF}(S_0, \mathbf{a}, \hbar(\mathbf{a})) \to ECC_{2g}(N)$, which was shown to be a quasi-isomorphism in [CGH2, CGH3]. Here we are using $ECC_{2g}(N)$ instead of $PFC_{2g}(N)$, but there is no substantial difference.

Observe that there is a discrepancy between the algebra and the geometry: the map Φ_{alg} which we are using here is not the map Φ, and we need to reconcile the two.

Proof. If $Z \in \widehat{CF}'(S_0, \mathbf{a}, \hbar(\mathbf{a}))$, then $\Phi^+(Z) = \Phi'(Z)$ by Theorem 5.8.3. We observed in Theorem 3.1.4 that $H(Z) = 0$. Moreover, $K(Z) = 0$ by Theorem 5.9.2 and thus $K^+(Z) = 0$. Hence

$$\Phi_{alg}\begin{pmatrix} Z \\ 0 \end{pmatrix} = \begin{pmatrix} \Phi^+(Z) \\ K^+(Z) \end{pmatrix} = \begin{pmatrix} \Phi'(Z) \\ 0 \end{pmatrix},$$

and the following diagram is commutative:

$$\begin{tikzcd}
\widehat{CF}'(S_0, \mathbf{a}, \hbar(\mathbf{a})) \arrow[r, "\Phi'"] \arrow[d, "i \circ j'"'] & ECC_{2g}(N) \arrow[d, "\mathfrak{i} \circ q_2"'] \\
C(U) \arrow[r, "\Phi_{alg}"] & C(U').
\end{tikzcd}$$

This gives rise to the following commutative diagram of homology groups:

$$\begin{tikzcd}
\widehat{HF}(S_0, \mathbf{a}, \hbar(\mathbf{a})) \arrow[r, "\Phi_*"] \arrow[d, "i_* \circ j"'] & ECH_{2g}(N) \arrow[d, "(\mathfrak{i} \circ q_2)_*"'] \\
H(C(U)) \arrow[r, "(\Phi_{alg})_*"] & H(C(U')).
\end{tikzcd}$$

Since j, i_*, Φ_*, $(q_2)_*$, and $\mathfrak{i}_*$ are isomorphisms by Lemma 6.2.1, Lemma 6.2.2, [CGH2, CGH3], Lemma 6.3.1, and Lemma 6.3.3, Φ_{alg} itself is a quasi-isomorphism. $\square$

Acknowledgements. We are indebted to Michael Hutchings for many helpful conversations and for our previous collaboration which was a catalyst for the present work. We also thank Tobias Ekholm, Dusa McDuff, Ivan Smith and Jean-Yves Welschinger for illuminating exchanges. Part of this work was done while KH and PG visited MSRI during the academic year 2009–2010. We are extremely grateful to MSRI and the organizers of the "Symplectic and Contact Geometry and Topology" and the "Homology Theories of Knots and Links" programs for their hospitality; this work probably would never have seen the light of day without the large amount of free time which was made possible by the visit to MSRI.

REFERENCES

[Bo1] F. Bourgeois, *A Morse Bott approach to contact homology*, from: "Symplectic and contact topology: interactions and perspectives (Toronto, ON/Montréal, QC, 2001)", Fields Inst. Comm. **35**, Amer. Math. Soc., Providence, RI (2003) 55–77.

[Bo2] F. Bourgeois, Ph.D. thesis.

[BEE] F. Bourgeois, T. Ekholm and Y. Eliashberg, *Effect of Legendrian surgery. With an appendix by Sheel Ganatra and Maksim Maydanskiy*, Geom. Topol. **16** (2012), 301-389.

[BEHWZ] F. Bourgeois, Y. Eliashberg, H. Hofer, K. Wysocki and E. Zehnder, *Compactness results in symplectic field theory*, Geom. Topol. **7** (2003), 799–888 (electronic).

[CGH1] V. Colin, P. Ghiggini and K. Honda, *Embedded contact homology and open book decompositions*, preprint 2010.

[CGH2] V. Colin, P. Ghiggini and K. Honda, *The equivalence of Heegaard Floer homology and embedded contact homology I*, preprint 2012.

[CGH3] V. Colin, P. Ghiggini and K. Honda, *The equivalence of Heegaard Floer homology and embedded contact homology II*, preprint 2012.

[CGHH] V. Colin, P. Ghiggini, K. Honda and M. Hutchings, *Sutures and contact homology I*, Geom. Topol. **15** (2011), 1749-1842.

[CHL] V. Colin, K. Honda and F. Laudenbach, *On the flux of pseudo-Anosov homeomorphisms*, Algebr. Geom. Topol. **8** (2008), 2147–2160.

[DS] S. Donaldson and I. Smith, *Lefschetz pencils and the canonical class for symplectic four-manifolds*, Topology **42** (2003), 743–785.

[Dr] D. Dragnev, *Fredholm theory and transversality for noncompact pseudoholomorphic maps in symplectizations*, Comm. Pure Appl. Math. **57** (2004), 726–763.

[EGH] Y. Eliashberg, A. Givental and H. Hofer, *Introduction to symplectic field theory*, GAFA 2000 (Tel Aviv, 1999), Geom. Funct. Anal. 2000, Special Volume, Part II, 560–673.

[Gi1] E. Giroux, *Convexité en topologie de contact*, Comment. Math. Helv. **66** (1991), 637–677.

[Gi2] E. Giroux, *Géométrie de contact: de la dimension trois vers les dimensions supérieures*, Proceedings of the International Congress of Mathematicians, Vol. II (Beijing, 2002), 405–414, Higher Ed. Press, Beijing, 2002.

[Gr] M. Gromov, *Pseudo holomorphic curves in symplectic manifolds*, Invent. Math. **82** (1985), 307–347.

[Ho1] H. Hofer, *Pseudoholomorphic curves in symplectizations with applications to the Weinstein conjecture in dimension three*, Invent. Math. **114** (1993), 515–563.

[Ho2] H. Hofer, *Holomorphic curves and dynamics in dimension three*, Symplectic geometry and topology (Park City, UT, 1997), 35–101, IAS/Park City Math. Ser., 7, Amer. Math. Soc., Providence, RI, 1999.

[HLS] H. Hofer, V. Lizan and J.-C. Sikorav, *On genericity for holomorphic curves in four-dimensional almost complex manifolds*, J. Geom. Anal. **7** (1997), 149–159.

[HWZ] H. Hofer, K. Wysocki and E. Zehnder, *Properties of pseudoholomorphic curves in symplectisations. I. Asymptotics*, Ann. Inst. H. Poincaré Anal. Non Linéaire **13** (1996), 337–379.

[HWZ2] H. Hofer, K. Wysocki and E. Zehnder, *Properties of pseudo-holomorphic curves in symplectizations. II. Embedding controls and algebraic invariants*, Geom. Funct. Anal. **5** (1995), 270–328.

[H] K. Honda, *On the classification of tight contact structures I*, Geom. Topol. **4** (2000) 309-368.

[HKM] K. Honda, W. Kazez and G. Matić, *On the contact class in Heegaard Floer homology*, J. Differential Geom. **83** (2009), 289–311.

[HKM2] K. Honda, W. Kazez and G. Matić, *Contact structures, sutured Floer homology, and TQFT*, preprint 2008. `ArXiv:0807.2431`.

[Hu] M. Hutchings, *An index inequality for embedded pseudoholomorphic curves in symplectizations*, J. Eur. Math. Soc. (JEMS) **4** (2002), 313–361.

[Hu2] M. Hutchings, *The embedded contact homology index revisited*, New perspectives and challenges in symplectic field theory, 263–297, CRM Proc. Lecture Notes, 49, AMS, 2009.

[HS1] M. Hutchings and M. Sullivan, *The periodic Floer homology of a Dehn twist*, Algebr. Geom. Topol. **5** (2005), 301–354.

[HS2] M. Hutchings and M. Sullivan, *Rounding corners of polygons and the embedded contact homology of* T^3, Geom. Topol. **10** (2006), 169–266 (electronic).

[HT1] M. Hutchings and C. Taubes, *Gluing pseudoholomorphic curves along branched covered cylinders I*, J. Symplectic Geom. **5** (2007), 43–137.

[HT2] M. Hutchings and C. Taubes, *Gluing pseudoholomorphic curves along branched covered cylinders II*, J. Symplectic Geom. **7** (2009), 29–133.

[HT3] M. Hutchings and C. Taubes, *Proof of the Arnold chord conjecture in three dimensions 1*, Math. Res. Lett. **18** (2011), 295-313.

[IP1] E. Ionel and T. Parker, *Relative Gromov-Witten invariants*, Ann. of Math. (2) **157** (2003), 45–96.

[IP2] E. Ionel and T. Parker, *The symplectic sum formula for Gromov-Witten invariants*, Ann. of Math. (2) **159** (2004), 935–1025.

[KM] M. Kontsevich and Y. Manin, *Gromov-Witten classes, quantum cohomology, and enumerative geometry*, Commun. Math. Phys. **164** (1994), 525–562.

[KLT1] C. Kutluhan, Y. Lee and C. Taubes, *HF=HM I: Heegaard Floer homology and Seiberg-Witten Floer homology*, preprint 2010. `ArXiv:1007.1979.`

[KLT2] C. Kutluhan, Y. Lee and C. Taubes, *HF=HM II: Reeb orbits and holomorphic curves for the ech/Heegaard-Floer correspondence*, preprint 2010. `ArXiv:1008.1595.`

[KLT3] C. Kutluhan, Y. Lee and C. Taubes, *HF=HM III: Holomorphic curves and the differential for the ech/Heegaard Floer correspondence*, preprint 2010. `ArXiv:1010.3456.`

[KLT4] C. Kutluhan, Y. Lee and C. Taubes, *HF=HM IV: The Seiberg-Witten Floer homology and ech correspondence*, preprint 2011. `ArXiv:1107.2297`.

[KLT5] C. Kutluhan, Y. Lee and C. Taubes, *HF=HM V: Seiberg-Witten-Floer homology and handle addition*, preprint 2012. `ArXiv:1204.0115`.

[LT] Y.-J. Lee and C. Taubes, *Periodic Floer homology and Seiberg-Witten Floer cohomology*, preprint 2009. `ArXiv:0906.0383.`

[Li] R. Lipshitz, *A cylindrical reformulation of Heegaard Floer homology*, Geom. Topol. **10** (2006), 955–1097.

[M1] D. McDuff, *The local behaviour of holomorphic curves in almost complex* 4*-manifolds*, J. Differential Geom. **34** (1991), 143–64.

[M2] D. McDuff, *Singularities and positivity of intersections of J-holomorphic curves. With an appendix by Gang Liu*, Progr. Math., 117, Holomorphic curves in symplectic geometry, 191-215, Birkhäuser, Basel, 1994.

[MS] D. McDuff and D. Salamon, *J-holomorphic curves and symplectic topology*, American Mathematical Society Colloquium Publications, 52. American Mathematical Society, Providence, RI, 2004.

[MW] M. Micallef and B. White, *The structure of branch points in minimal surfaces and in pseudo-holomorphic curves*, Ann. of Math. (2) **141** (1995), 35–85.

[OSz1] P. Ozsváth and Z. Szabó, *Holomorphic disks and topological invariants for closed three-manifolds*, Ann. of Math. (2) **159** (2004), 1027–1158.

[OSz2] P. Ozsváth and Z. Szabó, *Holomorphic disks and three-manifold invariants: properties and applications*, Ann. of Math. (2) **159** (2004), 1159–1245.

[Se1] P. Seidel, *A long exact sequence for symplectic Floer cohomology*, Topology **42** (2003), 1003–1063.

[Se2] P. Seidel, *Fukaya categories and Picard-Lefschetz theory.* Zurich Lectures in Advanced Mathematics. European Mathematical Society (EMS), Zürich, 2008.

[T1] C. Taubes, *The Seiberg-Witten equations and the Weinstein conjecture*, Geom. Topol. **11** (2007), 2117–2202.

[T2] C. Taubes, *Embedded contact homology and Seiberg-Witten Floer cohomology I–V*, Geom. Topol. **14** (2010) 2497–3000.

[T3] C. Taubes, *The structure of pseudoholomorphic subvarieties for a degenerate almost complex structure and symplectic form on* $S^1 \times B^3$, Geom. Topol. **2** (1998), 221–332.

[T4] C. Taubes, *Counting pseudo-holomorphic submanifolds in dimension* 4, Seiberg Witten and Gromov invariants for symplectic 4-manifolds, 99-161, First Int. Press Lect. Ser., 2, Int. Press, Somerville, MA, 2000.

[We1] C. Wendl, *Finite energy foliations on overtwisted contact manifolds*, Geom. Topol. **12** (2008) 531–616.

[We2] C. Wendl, *Finite energy foliations and surgery on transverse links*, Ph.D. thesis 2005.

[We3] C. Wendl, *Automatic transversality and orbifolds of punctured holomorphic curves in dimension four*, preprint 2008. `ArXiv:0802.3842`.

[We4] C. Wendl, *Non-exact symplectic cobordisms between contact* 3*-manifolds*, `ArXiv:10008.2456`

UNIVERSITÉ DE NANTES, 44322 NANTES, FRANCE
E-mail address: `vincent.colin@univ-nantes.fr`

UNIVERSITÉ DE NANTES, 44322 NANTES, FRANCE
E-mail address: `paolo.ghiggini@univ-nantes.fr`
URL: `http://www.math.sciences.univ-nantes.fr/˜Ghiggini`

UNIVERSITY OF SOUTHERN CALIFORNIA, LOS ANGELES, CA 90089
E-mail address: `khonda@usc.edu`
URL: `http://www-bcf.usc.edu/˜khonda`